



BlueCross BlueShield  
of Alabama

---

**Name of Blue Advantage Policy:**

**Bi-Ventricular Pacemakers (Cardiac Resynchronization Therapy)  
for the Treatment of Heart Failure**

Policy #: 055  
Category: Surgery

Latest Review Date: May 2021  
Policy Grade: A

---

**BACKGROUND:**

*Blue Advantage medical policy does not conflict with Local Coverage Determinations (LCDs), Local Medical Review Policies (LMRPs) or National Coverage Determinations (NCDs) or with coverage provisions in Medicare manuals, instructions or operational policy letters. In order to be covered by Blue Advantage the service shall be reasonable and necessary under Title XVIII of the Social Security Act, Section 1862(a)(1)(A). The service is considered reasonable and necessary if it is determined that the service is:*

1. *Safe and effective;*
2. *Not experimental or investigational\*;*
3. *Appropriate, including duration and frequency that is considered appropriate for the service, in terms of whether it is:*
  - *Furnished in accordance with accepted standards of medical practice for the diagnosis or treatment of the patient's condition or to improve the function of a malformed body member;*
  - *Furnished in a setting appropriate to the patient's medical needs and condition;*
  - *Ordered and furnished by qualified personnel;*
  - *One that meets, but does not exceed, the patient's medical need; and*
  - *At least as beneficial as an existing and available medically appropriate alternative.*

*\*Routine costs of qualifying clinical trial services with dates of service on or after September 19, 2000 which meet the requirements of the Clinical Trials NCD are considered reasonable and necessary by Medicare. Providers should bill **Original Medicare** for covered services that are related to **clinical trials** that meet Medicare requirements (Refer to Medicare National Coverage Determinations Manual, Chapter 1, Section 310 and Medicare Claims Processing Manual Chapter 32, Sections 69.0-69.11).*

## **POLICY:**

### **Effective for dates of service on or after June 8, 2018:**

**Blue Advantage will treat biventricular pacemakers, with or without an accompanying implantable cardiac defibrillator (i.e., a combined biventricular pacemaker/ICD) as a covered benefit for the treatment of heart failure (HF) when ALL of the following criteria are met:**

New York Heart Association (NYHA) Class III or IV

- Left ventricular ejection fraction  $\leq 35\%$ .
- Sinus rhythm
- Patients treated with guideline-directed medical therapy\*

**AND**

- Either QRS duration of  $\geq 120$  msec\*\* - or left bundle branch block

New York Heart Association (NYHA) Class II

- Left ventricular ejection fraction  $\leq 30\%$
- Sinus rhythm
- Patients treated with guideline-directed medical therapy\*

**AND**

- Either QRS duration of  $\geq 120$  msec\*\* or left bundle branch block

**Blue Advantage patients who do not meet the criteria outlined above, but have an indication for a ventricular pacemaker, biventricular pacemakers with or without an accompanying implantable cardiac defibrillator (i.e., a combined biventricular pacemaker/ICD) will meet Blue Advantage's medical criteria for coverage as an alternative to a right ventricular pacemaker in patients who meet ALL of the following criteria:**

- NYHA class I, II, III, or IV heart failure
- Left ventricular ejection fraction  $\leq 50\%$
- The presence of atrioventricular block with requirement for a high percentage of ventricular pacing\*\*\*
- Patients treated with guideline directed medical therapy\*\*

\*Guideline-directed medical therapy for heart failure is outlined in 2013 American College of Cardiology Foundation/American Heart Association guidelines for the management of heart failure.

\*\*The FDA-labeled indications for QRS duration vary by device. For some devices, FDA approval is based on QRS duration of  $\geq 130$  (e.g., InSync® device), while for others, it is based on QRS duration  $\geq 120$  msec (e.g., Guidant). These differences in QRS duration arise from differences in the eligibility criteria in the trials on which the FDA approval is based.

\*\*\*Atrioventricular block with a requirement for a high percentage of ventricular pacing is considered to be present when there is either:

- 3<sup>rd</sup> degree atrioventricular block; OR

- 2<sup>nd</sup> degree atrioventricular block or a PR interval of 300ms or more when paced at 100 beats per minute.

**Blue Advantage** will treat a **combined biventricular pacemaker and implantable cardiac defibrillator (ICD)** as a **covered benefit** for patients who meet criteria for **BOTH** a biventricular pacemaker and an ICD. **Please see CMS National Coverage Determination (NCD) 20.4 for criteria for the ICD.**

**Blue Advantage** will treat **biventricular pacemakers, with or without an accompanying implantable cardiac defibrillator**, as a **non-covered** benefit and as **investigational**.

- Treatment for patients with NYHA class I heart failure who do not meet the above criteria
- Including but not limited to the following:
  - atrial Fibrillation
  - unstable angina
  - myocardial infarction
  - prior coronary artery revascularization or angioplasty within the past 3 months

**Blue Advantage** will treat **intrathoracic fluid monitoring sensor as a component of a biventricular pacemaker** as a **non-covered benefit** and as **investigational**.

**Blue Advantage** will treat **triple-site (triventricular) CRT**, using an **additional pacing lead** as a **non-covered benefit** and as **investigational**.

**Blue Advantage** will treat **cardiac resynchronization therapy with wireless left ventricular endocardial pacing** as a **non-covered benefit** and as **investigational**.

*Blue Advantage does not approve or deny procedures, services, testing, or equipment for our members. Our decisions concern coverage only. The decision of whether or not to have a certain test, treatment or procedure is one made between the physician and his/her patient. Blue Advantage administers benefits based on the members' contract and medical policies. Physicians should always exercise their best medical judgment in providing the care they feel is most appropriate for their patients. Needed care should not be delayed or refused because of a coverage determination.*

## **DESCRIPTION OF PROCEDURE OR SERVICE:**

Cardiac resynchronization therapy (CRT), which consists of synchronized pacing of the left and right ventricles, is intended to treat patients with heart failure and dyssynchronous ventricular contractions. Treatment involves placement of a device that paces both ventricles and which coordinates ventricular pacing to maximize cardiac pumping function and left ventricular ejection fraction (LVEF).

## **Heart Failure**

It is estimated that 20%-30% of patients with heart failure (HF) have intraventricular conduction disorders resulting in a contraction pattern that is not coordinated and a wide QRS interval on the electrocardiogram (ECG). This abnormality appears to be associated with increased morbidity and mortality.

## **Treatment**

Biventricular pacemakers using 3 leads (1 in the right atrium, 1 endocardial in the right ventricle, 1 epicardial for the left ventricle), also known as CRT, have been investigated as a technique to coordinate the contraction of the ventricles, thus improving patients' hemodynamic status.

Originally developed CRT devices typically used two ventricular leads for biventricular pacing. Devices and implantation techniques have been developed to allow for multisite pacing, with the goal of improving CRT response. This may be accomplished in 1 of 2 ways: through the use of multiple leads within the coronary sinus (triventricular pacing) or through the use of multipolar left ventricular pacing leads, which can deliver pacing stimuli at multiple sites. Wireless left ventricular endocardial pacing is also being evaluated for patients who are not candidates for or do not respond to standard epicardial pacing leads.

## **KEY POINTS:**

The most recent literature review was updated through March 31, 2021.

### **Summary of Evidence**

For individuals who have NYHA class III/IV heart failure with left ventricular ejection fraction  $\leq$  35% who are in sinus rhythm, treated with guideline-directed medical therapy, and have either LBBB or QRS duration  $\geq$ 120ms, who receive cardiac resynchronization therapy (CRT), the evidence includes randomized controlled trials (RCTs) and systematic reviews of RCTs. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, hospitalizations, and treatment-related morbidity. There is a large body of clinical trial evidence supporting use of CRT in patients with NYHA class III or IV heart failure. The RCTs have consistently reported that CRT treatment reduces mortality, improves functional status, and improves quality of life for patients with NYHA class III or IV heart failure. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have NYHA class II heart failure with a left ventricular ejection fraction of 30% or less who are in sinus rhythm, treated with guideline-directed medical therapy, and have either LBBB or a QRS interval of 120 ms or more who receive CRT with or without defibrillator, the evidence includes RCTs and systematic reviews of RCTs. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, hospitalizations, and treatment-related morbidity. For patients with NYHA class II heart failure, at least 4 RCTs assessing CRT have been published. A mortality benefit was reported in 1 of the 4 trials, the Resynchronization-Defibrillation for Ambulatory Heart Failure Trial. None of the other 3 RCTs reported a mortality difference, but a subgroup analysis of the MADIT-CRT trial reported a mortality benefit for patients with LBBB. Among other outcome measures, hospitalizations for heart failure showed consistent reductions, but quality of life and functional status did not

improve The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have NYHA class I heart failure who receive CRT with or without defibrillator, the evidence includes RCTs and systematic reviews of RCTs. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, hospitalizations, and treatment-related morbidity. Few patients with NYHA class I heart failure have been included in RCTs. The MADIT-CRT trial included 265 patients with class I. While the treatment effect on death and hospitalization favored combined implantable cardiac defibrillator plus CRT devices vs implantable cardiac defibrillator alone for class I patients, the confidence interval was large and included a 25% to 30% increase in events. The evidence is insufficient to determine that the technology results in an improvement in net health outcomes.

For individuals who have NYHA class I, II, III or IV heart failure with left ventricular ejection fraction of 50% or less and the presence of atrioventricular block with requirement for a high percentage of ventricular pacing, treated with guideline-directed medical therapy, who receive CRT with or without defibrillator, the evidence includes RCTs and systematic reviews of RCTs. Relevant outcomes are OS, symptoms, functional outcomes, quality of life, hospitalizations, and treatment-related morbidity. For patients who have atrioventricular nodal block, some degree of left ventricular dysfunction, and who would not necessarily meet conventional criteria for CRT but would require ventricular pacing, a large RCT has demonstrated improvements in heart failure-related hospitalizations and urgent care visits among patients treated with CRT instead of RV pacing alone. For patients who require ventricular pacing but have no left ventricular dysfunction, results of a small RCT have suggested that biventricular pacing is associated with improved measures of cardiac function, but the trial was small and underpowered to detect differences in clinical outcomes. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with atrial fibrillation and heart failure who receive CRT, the evidence consists of 5 RCTs and a registry study. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, hospitalizations, and treatment-related morbidity. Results from RCTs have reported conflicting results, with 1 reporting improvements for patients with atrial fibrillation (AF) and others reporting no significant improvements. A registry study reported significant improvements in mortality and hospitalizations for patients with heart failure and AF treated with CRT plus defibrillator compared with ICD alone. The evidence is insufficient to determine that the technology results in an improvement in the health outcomes.

For individuals who have heart failure and atrioventricular (AV) nodal block who receive CRT, the evidence includes RCTs. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, hospitalizations, and treatment-related morbidity. One large RCT demonstrated that CRT led to reductions in heart failure-related hospitalizations and urgent care visits among patients with heart failure and AV block but who would not necessarily meet conventional criteria for CRT. For patients who require ventricular pacing but have no left ventricular dysfunction, results of a small RCT have suggested that biventricular pacing is associated with improved measures of cardiac function, but the trial was small and underpowered

to detect differences in clinical outcomes. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have heart failure who receive triple-site CRT, the evidence includes small RCTs and a meta-analysis that included nonrandomized studies. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, hospitalizations, and treatment-related morbidity. The available RCTs have reported improved outcomes on at least 1 measure of functional status or quality of life with triple-site CRT compared to conventional CRT. However, the trials are small and have methodologic limitations. In addition, outcomes reported differed across studies. Triple-site CRT was also associated with higher radiation exposure and a greater number of additional procedures postimplantation. Larger, high-quality RCTs are needed to better define the benefit-risk ratio for triple-site CRT compared to conventional CRT. The evidence is insufficient to determine that the technology results in an improvement in the net health outcomes.

For individuals who have heart failure who receive CRT combined with remote fluid monitoring, the evidence includes 3 RCTs. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, hospitalizations, and treatment-related morbidity. Three RCTs have reported no improvement in outcomes associated with remote fluid monitoring for patients with heart failure. The evidence is insufficient to determine that the technology results in an improvement in the net health outcomes.

**Practice Guidelines and Position Statements**

**American College of Cardiology, American Heart Association, and Heart Rhythm Society**

The ACC and American Heart Association and Heart Rhythm Society (2019) published joint guidelines on the evaluation and management of patients with bradycardia and cardiac conduction delay. These guidelines included the following recommendations on CRT:

Recommendation	COR	LOE
"In patients with atrioventricular block who have an indication for permanent pacing with a LVEF between 36% and 50% and are expected to require ventricular pacing more than 40% of the time, it is reasonable to choose pacing methods that maintain physiologic ventricular activation (e.g., cardiac resynchronization therapy [CRT] or His bundle pacing) over right ventricular pacing."	IIa	B- RSR
"In patients with atrioventricular block who have an indication for permanent pacing with a LVEF between 36% and 50% and are expected to require ventricular pacing less than 40% of the time, it is reasonable to choose right ventricular pacing over pacing methods that maintain physiologic ventricular activation (e.g., CRT or His bundle pacing)."	IIa	B-R

COR: class of recommendation; LOE: level of evidence; LVEF: left ventricular ejection fraction.

In 2013, the American College of Cardiology (ACC) Foundation and American Heart Association (AHA) Task Force on Practice Guidelines published guidelines for the management of heart failure. These guidelines make recommendations regarding cardiac resynchronization therapy (CRT) for heart failure (HF) that are in line with those made by the ACC/AHA and Heart Rhythm Society related to CRT for HF outlined below.

A focused update to 2008 guidelines for device-based treatment of cardiac rhythm abnormalities were published jointly by ACC/AHA/HRS in 2012. These guidelines included the following recommendations on CRT for heart failure:

Class I recommendations:

- CRT is indicated for patients who have LVEF less than or equal to 35%, sinus rhythm, LBBB with a QRS duration greater than or equal to 150 ms, and NYHA Class II, III, or ambulatory IV symptoms on guideline-directed medical therapy (GDMT). (Level of Evidence: A for NYHA Class III/IV; Level of Evidence: B for NYHA Class II)

Class IIa recommendations:

- CRT can be useful for patients who have LVEF less than or equal to 35%, sinus rhythm, LBBB with a QRS duration 120 to 149 ms, and NYHA Class II, III, or ambulatory IV symptoms on GDMT. (Level of Evidence: B)
- CRT can be useful for patients who have LVEF less than or equal to 35%, sinus rhythm, a non-LBBB pattern with a QRS duration greater than or equal to 150 ms, and NYHA Class III/ambulatory Class IV symptoms on GDMT. (Level of Evidence: A)
- CRT can be useful in patients with atrial fibrillation and LVEF less than or equal to 35% on GDMT if a) the patient requires ventricular pacing or otherwise meets CRT criteria and b) AV nodal ablation or pharmacologic rate control will allow near 100% ventricular pacing with CRT. (Level of Evidence: B)
- CRT can be useful for patients on GDMT who have LVEF less than or equal to 35% and are undergoing new or replacement device placement with anticipated requirement for significant (>40%) ventricular pacing. (Level of Evidence: C)

Class IIb recommendations:

- CRT may be considered for patients who have LVEF less than or equal to 30%, ischemic etiology of heart failure, sinus rhythm, LBBB with a QRS duration of greater than or equal to 150 ms, and NYHA class I symptoms on GDMT. (Level of Evidence: C)
- CRT may be considered for patients who have LVEF less than or equal to 35%, sinus rhythm, a non-LBBB pattern with QRS duration 120 to 149 ms, and NYHA Class III/ambulatory Class IV on GDMT. (Level of Evidence: B)
- CRT may be considered for patients who have LVEF less than or equal to 35%, sinus rhythm, a non-LBBB pattern with a QRS duration greater than or equal to 150 ms, and NYHA Class II symptoms on GDMT. (Level of Evidence: B)

Class III recommendations (no benefit)

- CRT is not recommended for patients with NYHA Class I or II symptoms and non-LBBB pattern with QRS duration less than 150 ms. (Level of Evidence: B)
- CRT is not indicated for patients whose comorbidities and/or frailty limit survival with good functional capacity to less than one year. (Level of Evidence: C)

### **Heart Failure Society of America**

The Heart Failure Society of America released comprehensive guidelines on the management of heart failure in 2010. The guidelines include the following recommendations related to the use of CRT:

- Biventricular pacing therapy is recommended for patients in sinus rhythm with a widened QRS interval ( $\geq 120$  ms) and severe LV systolic dysfunction (LVEF  $\leq 35\%$ ) who have persistent, moderate to severe HF (NYHA III) despite optimal medical therapy. (Level of Evidence: A).
- Biventricular pacing therapy may be considered for patients with atrial fibrillation with a widened QRS interval ( $\geq 120$  ms) and severe LV systolic dysfunction LVEF  $\leq 35\%$  who have persistent, moderate to severe HF (NYHA III) despite optimal medical therapy. (Level of Evidence: B).
- Selected ambulatory NYHA IV patients in sinus rhythm with QRS  $\geq 120$  ms and LV systolic dysfunction may be considered for biventricular pacing therapy. (Level of Evidence: B).
- Biventricular pacing therapy may be considered in patients with reduced LVEF and QRS  $\geq 150$  ms who have NYHA I or II HF symptoms. (Level of Evidence: B).
- In patients with reduced LVEF who require chronic pacing and in whom frequent ventricular pacing is expected, biventricular pacing may be considered. (Level of Evidence: C).

### **National Institute for Health and Care Excellence**

The National Institute for Health and Care Excellence's 2014 guidance provided recommendations on cardiac resynchronization therapy for heart failure. The recommendations for patients with LVEF  $\leq 35\%$  are as follows:

- NYHA class I-IV with QRS interval  $< 120$  ms: CRT not recommended
- NYHA class IV with QRS interval 120 to 149 ms and without LBBB: CRT-P recommended
- NYHA class II-III with QRS interval 120 to 149 ms and with LBBB: CRT-D recommended
- NYHA class III-IV with QRS interval 120 to 149 ms and with LBBB: CRT-P recommended
- NYHA class I-III with QRS interval  $\geq 150$  ms (with or without LBBB): CRT-D recommended
- NYHA class III-IV with QRS interval  $\geq 150$  ms (with or without LBBB): CRT-P recommended

## U.S. Preventive Services Task Force Recommendations

Not applicable

### KEY WORDS:

InSync®, Biventricular Pacemaker, biventricular pacing, congestive heart failure (CHF), pacemaker, cardiac resynchronization, CRT, cardiac resynchronization therapy, Viva™ Quad XT, Viva Quad S, Attain Performa®, Dynagen, Inogen, OptiVol™, Triple site CRT, Triventricular Pacemaker, intrathoracic fluid monitoring sensor; WiSE-CRT, EBR Systems

### APPROVED BY GOVERNING BODIES:

There are numerous CRT devices, combined implantable cardiac defibrillator (ICD) plus CRT devices (CRT-D), and combined CRT plus fluid monitoring devices. Some of the devices are discussed here. For example, in 2001, a stand-alone biventricular pacemaker (InSync® Biventricular Pacing System, Medtronic) received approval by U.S. Food and Drug Administration (FDA) through the premarket approval (PMA) process for the treatment of patients with New York Heart Association (NYHA) Class III or IV heart failure, on a stable pharmacologic regimen, who also have a QRS duration of 130 ms or longer and a left ventricular ejection fraction (LVEF) of 35% or less. Both Guidant (CONTAK CD® CRT-D System) and Medtronic (InSync® ICD Model 7272) have received FDA approval through the PMA process for combined cardiac resynchronization therapy defibrillators for patients at high risk of sudden cardiac death due to ventricular arrhythmias and who have NYHA Class III or IV heart failure with LVEF of 35% or less, QRS duration 130 ms or longer ( $\geq 120$  ms for the Guidant device), and remain symptomatic despite a stable, optimal heart failure drug therapy. In 2006, Biotronik Inc. received premarket approval through the FDA for its combined ICD/CRT device with ventricular pacing leads (Tupos LV/ATx CRT-D/Kronos LV-T CRT-D systems); in 2013, the company received FDA approval for updated ICD/CRT devices (Ilesto/Iforia series).

On the basis of the MADIT-CRT study, indications for three Guidant (Boston Scientific) CRT-defibrillator devices (Cognis®, Livian®, and Contak Renewal devices) were expanded to include patients with heart failure who receive stable optimal pharmacologic therapy for heart failure and who meet any one of the following classifications:

- Moderate-to-severe heart failure (NYHA class III or IV) with ejection fraction less than 35% and QRS duration greater than 120ms.
- Left bundle branch block with QRS greater than or equal to 130ms, ejection fraction less than 30%, and mild (NYHA class II) ischemic or nonischemic heart failure or asymptomatic (NYHA class I) ischemic heart failure.

In April 2014, FDA further expanded the indications for multiple Medtronic CRT devices to include patients with NYHA functional class I, II, or III heart failure, who have LVEF of 50% or less on stable, optimal heart failure medical therapy, if indicated, and have AV block that is expected to require a high percentage of ventricular pacing that cannot be managed with algorithms to minimize right ventricular pacing. The expanded indication was based on data from the BLOCK-HF study, a Medtronic-sponsored RCT to evaluate the use of CRT in patients

with NYHA class I, II, or III heart failure, LVEF  $\leq$ 50%, and AV block.

Several CRT devices incorporate a fourth lead, providing quadripolar pacing. The Medtronic VIVA™ Quad XT and the Viva Quad S incorporate a fourth lead, the Medtronic Attain Performa® left ventricular lead, which received clearance for marketing from FDA in August 2014. The Dynagen™ X4 and Inogen™ X4 devices incorporate a fourth lead. Other CRT devices with quadripolar leads have been approved for use outside of the United States (e.g. St. Jude Quartet™ left ventricular lead).

Multiple devices manufactured by Medtronic combine a CRT with the OptiVol™ monitoring system. For example, in 2005, the InSync Sentry® system received FDA approval through the supplemental premarket approval (PMA) process. This combined biventricular pacemaker/ICD is also equipped to monitor intrathoracic fluid levels using bioimpedance technology, referred to as OptiVol™ Fluid Status Monitoring. Bioimpedance measures, defined as the electrical resistance of tissue to flow of current, are performed many times per day using a vector from the right ventricular coil on the lead in the right side of the heart to the implanted pacemaker devices; changes in bioimpedance reflect intrathoracic fluid status and are evaluated based on a computer algorithm. For example, changes in a patient's daily average of intrathoracic bioimpedance can be monitored; differences in the daily average compared with a baseline are reported as the OptiVol Fluid Index. It has been proposed that these data may be used as an early warning system of cardiac decompensation or to provide additional feedback enabling a physician to further tailor medical therapy. (See medical policy #441 – Cardiac Hemodynamic Monitoring for the Management of Heart Failure in the Outpatient Setting for stand alone devices)

The WiSE-CRT (EBR Systems) provides CRT with a small wireless electrode that is implanted within the left ventricle and controlled by ultrasound. It has European CE approval and is being studied in a multicenter pivotal trial.

### **BENEFIT APPLICATION:**

Coverage is subject to member's specific benefits. Group specific policy will supersede this policy when applicable.

### **CURRENT CODING:**

#### **CPT codes:**

0515T	Insertion of wireless cardiac stimulator for left ventricular pacing, including device interrogation and programming, and imaging supervision and interpretation, when performed; complete system (includes electrode and generator [transmitter and battery]) (Effective 01/01/2019)
0516T	Insertion of wireless cardiac stimulator for left ventricular pacing, including device

	interrogation and programming, and imaging supervision and interpretation, when performed; electrode only (Effective 01/01/2019)
0517T	Insertion of wireless cardiac stimulator for left ventricular pacing, including device interrogation and programming, and imaging supervision and interpretation, when performed; pulse generator component(s) (battery and/or transmitter) only (Effective 01/01/2019)
0518T	Removal of only pulse generator component(s) (battery and/or transmitter) of wireless cardiac stimulator for left ventricular pacing (Effective 01/01/2019)
0519T	Removal and replacement of wireless cardiac stimulator for left ventricular pacing; pulse generator component(s) (battery and/or transmitter) (Effective 01/01/2019)
0520T	Removal and replacement of wireless cardiac stimulator for left ventricular pacing; pulse generator component(s) (battery and/or transmitter), including placement of a new electrode (Effective 01/01/2019)
0521T	Interrogation device evaluation (in person) with analysis, review and report, includes connection, recording, and disconnection per patient encounter, wireless cardiac stimulator for left ventricular pacing (Effective 01/01/2019)
0522T	Programming device evaluation (in person) with iterative adjustment of the implantable device to test the function of the device and select optimal permanent programmed values with analysis, including review and report, wireless cardiac stimulator for left ventricular pacing (Effective 01/01/2019)
33207	Insertion of new or replacement of permanent pacemaker with transvenous electrode(s); ventricular
33208	Insertion or replacement of permanent pacemaker with transvenous electrode(s); atrial and ventricular
33213	Insertion of pacemaker pulse generator only; with existing dual leads
33217	Insertion of 2 transvenous electrodes, permanent pacemaker or implantable defibrillator
33221	Insertion of pacemaker pulse generator only; with existing multiple leads
33224	Insertion of pacing electrode, cardiac venous system, for left ventricular pacing, with attachment to previously placed pacemaker or pacing cardioverter-defibrillator pulse generator (including revision of pocket, removal insertion and/or replacement of existing

	generator)
33225	Insertion of pacing electrode, cardiac venous system, for left ventricular pacing, at time of insertion of implantable defibrillator or pacemaker pulse generator (i.e., upgrade to dual chamber system) (List separately in addition to code for primary procedure
33226	Repositioning of previously implanted cardiac venous system (left ventricular) electrode (including removal, insertion and/or replacement of existing generator)
33228	Removal of permanent pacemaker pulse generator with replacement of pacemaker pulse generator; dual lead system
33229	Removal of permanent pacemaker pulse generator with replacement of pacemaker pulse generator; multiple lead system
33233	Removal of permanent pacemaker pulse generator only

## REFERENCES:

1. Abraham WT, Compton S, Haas G, et al. Intrathoracic impedance vs daily weight monitoring for predicting worsening heart failure events: results of the Fluid Accumulation Status Trial (FAST). *Congest Heart Fail.* Mar-Apr 2011; 17(2): 51-55.
2. Abraham WT, Fisher WG, Smith AL, et al. Cardiac resynchronization in chronic heart failure. *N Engl J Med.* Jun 13 2002; 346(24):1845-1853.
3. Abraham WT, Young JB, Leon AR, et al. Effects of cardiac resynchronization on disease progression in patients with left ventricular systolic dysfunction, an indication for an implantable cardioverter-defibrillator, and mildly symptomatic chronic heart failure. *Circulation* 2004; 110(18): 2864-2868.
4. Achilli A, Sassara M, et al. Effectiveness of cardiac resynchronization therapy in patients with echocardiographic evidence of mechanical dyssynchrony. *J Cardiovasc Med,* February 2008; 9(2): 131-136.
5. Adabag S, Roukoz H, Anand IS et al. Cardiac resynchronization therapy in patients with minimal heart failure: a systematic review and meta-analysis. *J Am Coll Cardiol* 2011; 58(9):935-41.
6. Ali-Hassan-Sayegh S, Mirhosseini SJ, Karimi-Bondarabadi AA, et al. Cardiac resynchronization therapy in patients with mild heart failure is a reversal therapy. *Indian Heart J.* Jan - Feb 2017; 69(1):112-118.
7. Al-Majed NS, McAlister FA, Bakal JA et al. Meta-analysis: cardiac resynchronization therapy for patients with less symptomatic heart failure. *Ann Intern Med* 2011. [ePub ahead of print]

8. Anselme F, Bordachar P, Pasquie JL, et al. Safety, feasibility, and outcome results of cardiac resynchronization with triple-site ventricular stimulation compared to conventional cardiac resynchronization. *Heart Rhythm*. Jan 2016;13(1):183-189.
9. Auricchio A, Stellbrink C, Butter C, et al. Clinical efficacy of cardiac resynchronization therapy using left ventricular pacing in heart failure patients stratified by severity of ventricular conduction delay. *J Am Coll Cardiol*. Dec 17 2003; 42(12):2109-2116.
10. Auricchio A, Stellbrink C, Sack S, et al. Long-term clinical effect of hemodynamically optimized cardiac resynchronization therapy in patients with heart failure and ventricular conduction delay. *J Am Coll Cardiol*. Jun 19 2002; 39(12):2026-2033.
11. Bakker Patricia F, Meijburg Huub W, et al. Biventricular pacing in end-stage heart failure improves functional capacity and left ventricular function. *Journal of Interventional Cardiac Electrophysiology* 4, 2000, pp. 395-404.
12. Bencardino G, Di Monaco A, Russo E, et al. Outcome of Patients Treated by Cardiac Resynchronization Therapy Using a Quadripolar Left Ventricular Lead. *Circ J*. Feb 25 2016; 80(3):613-618.
13. Bertoldi EG, Polanczyk CA, Cunha V et al. Mortality reduction of cardiac resynchronization and implantable cardioverter-defibrillator therapy in heart failure: an updated meta-analysis. Does recent evidence change the standard of care? *J Card Fail* 2011; 17(10):860-6.
14. Beshai JF, Grimm RA, Nagueh SF et al. Cardiac-resynchronization therapy in heart failure with narrow QRS complexes. *N Engl J Med* 2007; 357(24):2461-71.
15. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Cardiac resynchronization therapy for mild congestive heart failure. *TEC Assessments* 2009; Volume 24, Tab 8.
16. Bohm M, Drexler H, Oswald H, et al. Fluid status telemedicine alerts for heart failure: a randomized controlled trial. *Eur Heart J*. Nov 01 2016; 37(41):3154-3163.
17. Boriani G, Kranig W, Donal E, et al. A randomized double-blind comparison of biventricular versus left ventricular stimulation for cardiac resynchronization therapy: the Biventricular versus Left Univentricular Pacing with ICD Back-up in Heart Failure Patients (B-LEFT HF) trial. *Am Heart J*. Jun 2010; 159(6):1052-1058.e1051.
18. Brachmann J, Bohm M, Rybak K et al. Fluid status monitoring with a wireless network to reduce cardiovascular-related hospitalizations and mortality in heart failure: rationale and design of the OptiLink HF Study (Optimization of Heart Failure Management using OptiVol Fluid Status Monitoring and CareLink). *Eur J Heart Fail* 2011; 13(7):796-804.
19. Brignole M, Auricchio A, Baron-Esquivias G, et al. 2013 ESC Guidelines on cardiac pacing and cardiac resynchronization therapy: the Task Force on cardiac pacing and resynchronization therapy of the European Society of Cardiology (ESC). Developed in collaboration with the European Heart Rhythm Association (EHRA). *Eur Heart J*. Aug 2013;34(29):2281-2329.
20. Brignole M, Botto G, Mont L et al. Cardiac resynchronization therapy in patients undergoing atrioventricular junction ablation for permanent atrial fibrillation: a randomized trial. *Eur Heart J* 2011; 32(19):2420-9.

21. Brignole M, Pokushalov E, Pentimalli F, et al. A randomized controlled trial of atrioventricular junction ablation and cardiac resynchronization therapy in patients with permanent atrial fibrillation and narrow QRS. *Eur Heart J*. Dec 01 2018; 39(45): 3999-4008.
22. Bristow MR, Saxon LA, Boehmer J et al. Cardiac-resynchronization therapy with or without an implantable defibrillator in advanced chronic heart failure. *N Engl J Med* 2004; 350(21):2140-50.
23. Bryant AR, Wilton SB, Lai MP, et al. Association between QRS duration and outcome with cardiac resynchronization therapy: A systematic review and meta-analysis. *J Electrocardiol*. Mar 2013; 46(2): 147-155.
24. Cazeau Serge, Leclercq Christophe, et al. Effects of multisite biventricular pacing in patients with heart failure and intraventricular conduction delay. *New England Journal of Medicine*, March 2001, Vol. 344, pp. 873-880.
25. Chan JY, Fang F, Zhang Q, et al. Biventricular pacing is superior to right ventricular pacing in bradycardia patients with preserved systolic function: 2-year results of the PACE trial. *Eur Heart J*. Oct 2011; 32(20):2533- 2540.
26. Chen S, Ling Z, Kiuchi MG, et al. The efficacy and safety of cardiac resynchronization therapy combined with implantable cardioverter defibrillator for heart failure: a meta-analysis of 5674 patients. *Europace*. Jul 2013; 15(7): 992-1001.
27. Chung ES, Leon AR, Tavazzi L et al. Results of the Predictors of Response to CRT (PROSPECT) trial. *Circulation* 2008; 117(20):2608-16.
28. Cleland JG, Daubert JC, Erdmann E, et al. The effect of cardiac resynchronization on morbidity and mortality in heart failure. *N Engl J Med*. Apr 14 2005; 352(15):1539-1549.
29. Conraads VM, Tavazzi L, Santini M et al. Sensitivity and positive predictive value of implantable intrathoracic impedance monitoring as a predictor of heart failure hospitalizations: the SENSE-HF trial. *European Heart Journal* 2011.
30. Curtis AB, Worley SJ, Adamson PB, et al. Biventricular pacing for atrioventricular block and systolic dysfunction. *N Engl J med*. Apr 25 2013; 368 (17): 1585-1593.
31. Curtis AB, Worley SJ, Chung ES, et al. Improvement in clinical outcomes with biventricular versus right ventricular pacing: The BLOCK HF Study. *J Am Coll Cardiol*. May 10 2016; 67(18):2148-2157.
32. Diab IG, Hunter RJ, Kamdar R et al. Does ventricular dyssynchrony on echocardiography predict response to cardiac resynchronisation therapy? A randomised controlled study. *Heart* 2011; 97(17):1410-6.
33. Domenichini G, Rahneva T, Diab IG, et al. The lung impedance monitoring in treatment of chronic heart failure (the LIMIT-CHF study). *Europace*. Mar 2016; 18(3):428-435.
34. Doshi RN, Daoud EG, Fellows C, et al. Left ventricular-based cardiac stimulation post AV nodal ablation evaluation (the PAVE study). *J Cardiovasc Electrophysiol*. Nov 2005; 16(11):1160-1165.
35. Epstein AE, DiMarco JP, Ellenbogen KA et al. ACC/AHA/HRS 2008 Guidelines for Device-Based Therapy of Cardiac Rhythm Abnormalities: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines

- (Writing Committee to Revise the ACC/AHA/NASPE 2002 Guideline Update for Implantation of Cardiac Pacemakers and Antiarrhythmia Devices): developed in collaboration with the American Association for Thoracic Surgery and Society of Thoracic Surgeons. *Circulation* 2008; 117(21):e350-408.
36. Eriksson H. Heart failure: A growing public health problem. *Journal of Internal Medicine*, 1995; 237:135-141.
  37. European Society of C, European Heart Rhythm A, Brignole M et al. 2013 ESC guidelines on cardiac pacing and cardiac resynchronization therapy: the task force on cardiac pacing and resynchronization therapy of the European Society of Cardiology (ESC). Developed in collaboration with the European Heart Rhythm Association (EHRA). *Europace* 2013; 15(8):1070-118.
  38. Ezekowitz JA, Rowe BH, Dryden DM, et al. Systematic review: implantable cardioverter defibrillators for adults with left ventricular systolic dysfunction. *Ann Intern Med*. Aug 21 2007;147(4):251-262.
  39. FDA. Approval Order: Biotronic PMA P050023. 2013. Available online at: [www.accessdata.fda.gov/cdrh\\_docs/pdf5/P050023S058A.pdf](http://www.accessdata.fda.gov/cdrh_docs/pdf5/P050023S058A.pdf). Last accessed. Accessed April 8, 2021.
  40. FDA. Summary of Safety and Effectiveness Data: Tupos LV/ATx CRT-D, Kronos LV-T CRT-D. 2006. Available online at: [www.accessdata.fda.gov/cdrh\\_docs/pdf5/P050023b.pdf](http://www.accessdata.fda.gov/cdrh_docs/pdf5/P050023b.pdf). Accessed April 20, 2018.
  41. FDA. Summary of Safety and Effectiveness Data, PMA P030054. 2005; [www.accessdata.fda.gov/cdrh\\_docs/pdf3/P030054b.pdf](http://www.accessdata.fda.gov/cdrh_docs/pdf3/P030054b.pdf). Accessed April 20, 2018.
  42. FDA. Summary of Safety and Effectiveness Data, PMA P030035. 2005; [www.fda.gov/ohrms/dockets/dockets/05m0289/05m-0289-aav0001-03-SSED-vol1.pdf](http://www.fda.gov/ohrms/dockets/dockets/05m0289/05m-0289-aav0001-03-SSED-vol1.pdf). Accessed March 31, 2016
  43. Food and Drug Administration. Summary of Safety and Effectiveness Data: Cardiac Resynchronization Therapy Defibrillator (CRT-D). 2010; [www.accessdata.fda.gov/cdrh\\_docs/pdf/P010012S230b.pdf](http://www.accessdata.fda.gov/cdrh_docs/pdf/P010012S230b.pdf). Accessed April 8, 2021.
  44. Food and Drug Administration. Summary of Safety and Effectiveness Data: Cardiac Resynchronization Therapy - Pacemaker (CRT-P) System. PMA P030035. 2005; [https://www.accessdata.fda.gov/cdrh\\_docs/pdf3/p030035s003b.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf3/p030035s003b.pdf). Accessed April 20, 2018.
  45. Foley PW, Patel K, Irwin N, et al. Cardiac resynchronisation therapy in patients with heart failure and a normal QRS duration: the RESPOND study. *Heart*. Jul 2011; 97(13):1041-1047.
  46. Foreman B, Fishel RS, Odryzynski NI et al. Intra-thoracic impedance: a surrogate measure of thoracic fluid – Fluid Accumulation Status Trial (FAST). *J Card Fail* 2004; 10(suppl):251.
  47. Friedman DJ, Bao H, Spatz ES, et al. Association between a prolonged pr interval and outcomes of cardiac resynchronization therapy: a report from the National Cardiovascular Data Registry. *Circulation*. Nov 22 2016; 134(21):1617-1628.

48. Ganesan AN, Brooks AG, Roberts-Thomson KC et al. Role of AV nodal ablation in cardiac resynchronization in patients with coexistent atrial fibrillation and heart failure: a systematic review. *J Am Coll Cardiol* 2012; 59(8):719-26.
49. Garrigue S, Bordachar P, Reuter S, et al. Comparison of permanent left ventricular and biventricular pacing in patients with heart failure and chronic atrial fibrillation: prospective haemodynamic study. *Heart*. Jun 2002; 87(6):529-534.
50. Gasparini M, Bocchiardo M, Lunati M, et al. Comparison of 1-year effects of left ventricular and biventricular pacing in patients with heart failure who have ventricular arrhythmias and left bundle-branch block: the Bi vs Left Ventricular Pacing: an International Pilot Evaluation on Heart Failure Patients with Ventricular Arrhythmias (BELIEVE) multicenter prospective randomized pilot study. *Am Heart J*. Jul 2006; 152(1):155.e151-157.
51. Gasparini M, Auricchio A, Regoli F et al. Four-year efficacy of cardiac resynchronization therapy on exercise tolerance and disease progression: the importance of performing atrioventricular junction ablation in patients with atrial fibrillation. *J Am Coll Cardiol* 2006; 48(4):734-43.
52. Gillis AM, Kerr CR, Philippon F. et al. Impact of cardiac resynchronization therapy on hospitalizations in the resynchronization-defibrillation for ambulatory heart failure trial. *Circulation* May 20 2014; 129(20): 2021-2030.
53. Gold MR, Padhiar A, Mealing S, et al. Long-Term Extrapolation of Clinical Benefits Among Patients With Mild Heart Failure Receiving Cardiac Resynchronization Therapy: Analysis of the 5-Year Follow-Up From the REVERSE Study. *JACC Heart Fail*. Sep 2015; 3(9):691-700.
54. Goldenberg I, Hall WJ, Beck CA et al. Reduction of the risk of recurring heart failure events with cardiac resynchronization therapy: MADIT-CRT (Multicenter Automatic Defibrillator Implantation Trial With Cardiac Resynchronization Therapy). *J Am Coll Cardiol* 2011; 58(7):729-37.
55. Goldenberg I, Kutyifa V, Klein HU, et al. Survival with cardiac-resynchronization therapy in mild heart failure. *N Engl J Med*. May 1 2014; 370 (18): 1694-1701.
56. Gorcsan J III, Abraham T, Agler DA, et al. Echocardiography for cardiac resynchronization therapy: Recommendations for performance and reporting—a report from the American Society of Echocardiography Dyssynchrony Writing Group endorsed by the Heart Rhythm Society. *J Am Soc Echocardiogr*, March 2008; 21(3): 191-213.
57. Gula LJ, Wells GA, Yee R, et al. A novel algorithm to assess risk of heart failure exacerbation using ICD diagnostics: validation from RAFT. *Heart Rhythm* Sep 2014; 11(9): 1626-1631.
58. Hawkins NM, Petrie MC, MacDonald MR et al. Selecting patients for cardiac resynchronization therapy: electrical or mechanical dyssynchrony? *Eur Heart J* 2006; 27(11):1270-81.
59. Healey JS, Hohnloser SH, Exner DV et al. Cardiac resynchronization therapy in patients with permanent atrial fibrillation: results from the resynchronization for ambulatory heart failure trial (RAFT). *Circ Heart Fail*. Sep 1 2012; 5(5): 566-570.

60. Heart Failure Society of America, Lindenfeld J, Albert NM et al. HFSA 2010 Comprehensive Heart Failure Practice Guideline. *Journal of Cardiac Failure* 2010; 16(6):e1-194.
61. Higgins SL, Hummel JD, Niazi IK, et al. Cardiac resynchronization therapy for the treatment of heart failure in patients with intraventricular conduction delay and malignant ventricular tachyarrhythmias. *J Am Coll Cardiol*. Oct 15 2003; 42(8):1454-1459.
62. Hosseini SM, Moazzami K, Rozen G, et al. Utilization and in-hospital complications of cardiac resynchronization therapy: trends in the United States from 2003 to 2013. *Eur Heart J*. Mar 13 2017.
63. Hunt SA. ACC/AHA 2005 guideline update for the diagnosis and management of chronic heart failure in the adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to update the 2001 Guidelines for the Evaluation and Management of Heart Failure). *J Am Coll Cardiol* 2005; 46(6):e1-82.
64. Innes Donald, et al. VDD pacing at short atrioventricular intervals does not improve cardiac output in patients with dilated heart failure. *PACE*, May 1994, Vol. 17, Part I, pp. 956-965.
65. IOM (Institute of Medicine). 2011. *Clinical Practice Guidelines We Can Trust*. Washington, DC: The National Academies Press.
66. Jessup M. MADIT-CRT—Breathhtaking or time to catch our breath? *NEJM*, October 2009; 361(14): 1394-1338.
67. Kalscheur MM, Saxon LA, Lee BK, et al. Outcomes of cardiac resynchronization therapy in patients with intermittent atrial fibrillation or atrial flutter in the COMPANION trial. *Heart Rhythm*. Mar 17 2017.
68. Kang SH, Oh IY, Kang DY et al. Cardiac resynchronization therapy and QRS duration: systematic review, meta-analysis, and meta-regression. *J Korean Med Sci*. Jan 2015; 30(1): 24-33.
69. Kannel William B and Belanger Albert J. Epidemiology of heart failure. *American Heart Journal*, 1991; 121:951-957.
70. Kerwin Walter F, et al. Ventricular contraction abnormalities in dilated cardiomyopathy: Effect of Biventricular pacing to correct interventricular dyssynchrony. *Journal of American College of Cardiology*, 2000, Vol. 35, No. 5, pp. 1221-1227.
71. Khazanie P, Greiner MA, Al-Khatib SM, et al. Comparative effectiveness of cardiac resynchronization therapy among patients with heart failure and atrial fibrillation: findings from the National Cardiovascular Data Registry's Implantable Cardioverter-Defibrillator Registry. *Circ Heart Fail*. Jun 2016; 9(6).
72. Kindermann M, Hennen B, Jung J, et al. Biventricular versus conventional right ventricular stimulation for patients with standard pacing indication and left ventricular dysfunction: the Homburg Biventricular Pacing Evaluation (HOBIPACE). *J Am Coll Cardiol*. May 16 2006; 47(10):1927-1937.
73. Kusumoto FM, Schoenfeld MH, Barrett C, et al. 2018 ACC/AHA/HRS Guideline on the Evaluation and Management of Patients With Bradycardia and Cardiac Conduction Delay: Executive Summary: A Report of the American College of Cardiology/American

Heart Association Task Force on Clinical Practice Guidelines, and the Heart Rhythm Society. *J Am Coll Cardiol.* Aug 20 2019; 74(7): 932-987.

74. Kutyifa V, Stockburger M, Daubert JP, et al. PR interval identifies clinical response in patients with non-left bundle branch block: a multicenter automatic defibrillator implantation trial-cardiac resynchronization therapy substudy. *Circ Arrhythm Electrophysiol.* Aug 2014; 7(4): 645-651.
75. Leclercq C, Gadler, Kranig W, et al. A randomized comparison of triple-site versus dual-site ventricular stimulation in patients with congestive heart failure. *JACC*, April 2008; 51(15): 1455-1462.
76. Leclercq C, Victor F, et al. Comparative effects of permanent biventricular pacing for refractory heart failure in patients with stable sinus rhythm or chronic atrial fibrillation. *Journal American College of Cardiology*, May 2000, Vol 85, pp. 1154-56.
77. Leclercq Christophe, Cazeau Serge, et al. Acute hemodynamic effects of biventricular DDD pacing in patients with end-stage heart failure. *Journal American College of Cardiology*, 1998; 32: 1825-31.
78. Leclercq C, Cazeau S, Lellouche D, et al. Upgrading from single chamber right ventricular to biventricular pacing in permanently paced patients with worsening heart failure: The RD-CHF Study. *Pacing Clin Electrophysiol.* Jan 2007; 30 Suppl 1:S23-30.
79. Lenarczyk R, Kowalski O, Sredniawa B et al. Implantation feasibility, procedure-related adverse events and lead performance during 1-year follow-up in patients undergoing triple-site cardiac resynchronization therapy: a substudy of TRUST CRT randomized trial. *J Cardiovasc Electrophysiol* 2012; 23(11):1228-36.
80. Lin J, Buhr KA, Kipp R. Effect of PR interval on outcomes following cardiac resynchronization therapy: a secondary analysis of the COMPANION trial. *J Cardiovasc Electrophysiol.* Feb 2017; 28(2):185-191.
81. Linde C, Abraham WT, Gold MR, et al. Randomized trial of cardiac resynchronization in mildly symptomatic heart failure patients and in asymptomatic patients with left ventricular dysfunction and previous heart failure symptoms. *JACC* 2008; 52(23): 1834-1843.
82. Linde C, Gadler F, et al. Results of atrioventricular synchronous pacing with optimized delay in patients with severe congestive heart failure. *Journal American College of Cardiology*, 1995; 75: 919-923.
83. Linde C, Gold MR, Abraham WT et al. Long term impact of cardiac resynchronization therapy in mild heart failure: 5 year results from the resynchronization reverses remodeling in systolic left ventricular dysfunction (REVERSE) study. *Eur Heart J.* Sep 2013; 34(33): 2592-2599.
84. Lindenfeld J, Albert NM, Boehmer JP, et al. HFSA 2010 Comprehensive Heart Failure Practice Guideline. *J Card Fail.* Jun 2010; 16(6): e1-194.
85. Lozano I, Bocchiardo M, Ahtelik M, et al. Impact of biventricular pacing on mortality in a randomized crossover study of patients with heart failure and ventricular arrhythmias. *Pacing Clin Electrophysiol.* Nov 2000; 23(11 Pt 2):1711-1712.

86. Luthje L, Vollmann D, Seegers J, et al. A randomized study of remote monitoring and fluid monitoring for the management of patients with implanted cardiac arrhythmia devices. *Europace*. Aug 2015; 17(8):1276-1281.
87. Martinelli Filho M, de Siqueira SF, Costa R, et al. Conventional versus biventricular pacing in heart failure and bradyarrhythmia: the COMBAT study. *J Card Fail*. Apr 2010; 16(4):293-300.
88. McAlister FA, Ezekowitz J, Hooton N, et al. Cardiac resynchronization therapy for patients with left ventricular systolic dysfunction: a systematic review. *Jama*. Jun 13 2007; 297(22):2502-2514.
89. McAlister FA, Ezekowitz JA, Wiebe N et al. Systematic review: cardiac resynchronization in patients with symptomatic heart failure. *Ann Intern Med* 2004; 141(5):381-90.
90. Molhoek Sander G, Bax Jeroen J, et al. Comparison of benefits from cardiac resynchronization therapy in patients with ischemic cardiomyopathy versus idiopathic dilated cardiomyopathy. *The American Journal of Cardiology*, April 2004, Vol. 93, No. 7.
91. Moss AJ, Hall J, Cannom DS, et al. Cardiac-resynchronization therapy for the prevention of heart failure events. *N Engl J Med* 2009; 361:1329-1338.
92. Muto C, Solimene F, Gallo P et al. A randomized study of cardiac resynchronization therapy defibrillator versus dual-chamber implantable cardioverter-defibrillator in ischemic cardiomyopathy with narrow QRS: the NARROW-CRT study. *Circ Arrhythm Electrophysiol* 2013; 6(3):538-45.
93. National Institute for Health and Care Excellence (NICE). Implantable cardioverter defibrillators and cardiac resynchronisation therapy for arrhythmias and heart failure, technology appraisal guidance [TA314]. 2014; [www.nice.org.uk/guidance/ta314](http://www.nice.org.uk/guidance/ta314). Accessed April 5, 2021.
94. Nery PB, Ha AC, Keren A et al. Cardiac resynchronization therapy in patients with left ventricular systolic dysfunction and right bundle branch block: a systematic review. *Heart Rhythm* 2011; 8(7):1083-7.
95. Nishimura Rick A, Hayes David L, et al. Mechanism of hemodynamic improvement by dual-chamber pacing for severe left ventricular dysfunction: An acute doppler and catheterization hemodynamic study. *J Am Coll Cardiol*, 1995; 25: 281-8.
96. Ogano M, Iwasaki YK, Tanabe J et al. Antiarrhythmic effect of cardiac resynchronization therapy with triple-site biventricular stimulation. *Europace* 2013; 15(10):1491-8.
97. Pappone C, Calovic Z, Vicedomini G, et al. Improving cardiac resynchronization therapy response with multipoint left ventricular pacing: Twelve-month follow-up study. *Heart Rhythm*. Jun 2015; 12(6):1250-1258.
98. Perego GB, Landolina M, Vergara G, et al. Implantable CRT device diagnostics identify patients with increased risk for heart failure hospitalization. *J Interv Card Electrophysiol*, December 2008; 23(3): 235-242.
99. Peterson PN, Greiner MA, Qualls LG et al. QRS duration, bundle-branch block morphology, and outcomes among older patients with heart failure receiving cardiac resynchronization therapy. *JAMA* 2013; 310(6):617-26.

100. Piccirillo G, Magri D, di Carlo S, et al. Influence of cardiac-resynchronization therapy on heart rate and blood pressure variability: 1-year follow-up. *Eur J Heart Fail.* Nov 2006; 8(7):716-722.
101. Piepoli MF, Villani GQ, Corra U, et al. Time course of effects of cardiac resynchronization therapy in chronic heart failure: benefits in patients with preserved exercise capacity. *Pacing Clin Electrophysiol.* Jun 2008; 31(6):701-708.
102. Pinter A, Mangat I, Korley V, et al. Assessment of resynchronization therapy on functional status and quality of life in patients requiring an implantable defibrillator. *Pacing Clin Electrophysiol.* Dec 2009; 32(12):1509-1519.
103. Rao RK, Kumar UN, Schafer J, et al. Reduced ventricular volumes and improved systolic function with cardiac resynchronization therapy: a randomized trial comparing simultaneous biventricular pacing, sequential biventricular pacing, and left ventricular pacing. *Circulation.* Apr 24 2007; 115(16):2136-2144.
104. Rogers DP, Lambiase PD, Lowe MD et al. A randomized double-blind crossover trial of triventricular versus biventricular pacing in heart failure. *Eur J Heart Fail* 2012; 14(5):495-505.
105. Ruschitzka F, Abraham WT, Singh JP et al. Cardiac-resynchronization therapy in heart failure with a narrow QRS complex. *New England Journal of Medicine* 2013; 369(15):1395-405.
106. Santangeli P, Di Biase L, Pelargonio G, et al. Cardiac resynchronization therapy in patients with mild heart failure: a systematic review and meta-analysis. *J Interv Card Electrophysiol.* Nov 2011; 32(2):125-135.
107. Sekiguchi Y, Tada H, Yoshida K et al. Significant increase in the incidence of ventricular arrhythmic events after an intrathoracic impedance change measured with a cardiac resynchronization therapy defibrillator. *Circ J* 2011; 75(11):2614-20.
108. Shah RM, Patel D, Molnar J, et al. Cardiac-resynchronization therapy in patients with systolic heart failure and QRS interval  $\leq$  130ms: insights from a meta-analysis. *Europace* Feb 2015; 17(2): 267-273.
109. Shamim Waqar, et al. Intraventricular conduction delay: A prognostic marker in chronic heart failure. *International Journal of Cardiology*, 1999; 70: 171-178.
110. Sipahi I, Carrigan TP, Rowland DY et al. Impact of QRS duration on clinical event reduction with cardiac resynchronization therapy: meta-analysis of randomized controlled trials. *Arch Intern Med* 2011; 171(16):1454-62.
111. Sipahi I, Chou JC, Hyden M et al. Effect of QRS morphology on clinical event reduction with cardiac resynchronization therapy: meta-analysis of randomized controlled trials. *Am Heart J.* Feb 2012; 163(2): 260-267.
112. Stavrakis S, Lazzara R, Thadani U. The benefit of cardiac resynchronization therapy and QRS duration: a meta-analysis. *J Cardiovasc Electrophysiol.* Feb 2012; 23(2): 163-168.
113. Strickberger SA, Conti J, Daoud EG et al. Patient selection for cardiac resynchronization therapy: from the Council on Clinical Cardiology Subcommittee on Electrocardiography and Arrhythmias and the Quality of Care and Outcomes Research Interdisciplinary Working Group, in collaboration with the Heart Rhythm Society. *Circulation* 2005; 111(16):2146-50.

114. Stockburger M, Moss AJ, Klein HU, et al. Sustained clinical benefit of cardiac resynchronization therapy in non LBBB patients with prolonged PR-interval: MADIT-CRT long-term follow-up. *Clin Res Cardiol*. Nov 2016; 105(11):944-952.
115. Sun WP, Li CL, Guo JC, et al. Long-term efficacy of implantable cardiac resynchronization therapy plus defibrillator for primary prevention of sudden cardiac death in patients with mild heart failure: an updated metaanalysis. *Heart Fail Rev*. Jul 2016; 21(4):447-453.
116. Tang AS, Wells GA, Talajic M et al. Cardiac-resynchronization therapy for mild-to-moderate heart failure. *N Engl J Med* 2010; 363(25):2385-95.
117. Thibault B, Ducharme A, Harel F, et al. Left ventricular versus simultaneous biventricular pacing in patients with heart failure and a QRS complex  $\geq 120$  milliseconds. *Circulation*. Dec 20 2011; 124(25):2874-2881.
118. Thibault B, harel F, Ducharme A, et al. Cardiac resynchronization therapy in patients with heart failure and a QRS complex  $< 120$  milliseconds: the evaluation of resynchronization therapy for heart failure (LESSER-EARTH) trial. *Circulation*. Feb 26 2013; 127(8): 873-881.
119. Tracy CM, Epstein AE, Darbar D et al. 2012 ACCF/AHA/HRS Focused Update of the 2008 Guidelines for Device-Based Therapy of Cardiac Rhythm Abnormalities: A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines and the Heart Rhythm Society. *Circulation* 2012; 126(14):1784-800.
120. Tu R, Zhong G, Zeng Z et al. Cardiac resynchronization therapy in patients with mild heart failure: a systematic review and meta-analysis of randomized controlled trials. *Cardiovasc Drugs Ther* 2011; 25(4):331-40.
121. van Geldorp IE, Vernooy K, Delhaas T, et al. Beneficial effects of biventricular pacing in chronically right ventricular paced patients with mild cardiomyopathy. *Europace*. Feb 2010; 12(2):223-229.
122. van Rees JB, de Bie MK, Thijssen J et al. Implantation-related complications of implantable cardioverter-defibrillators and cardiac resynchronization therapy devices: a systematic review of randomized clinical trials. *J Am Coll Cardiol* 2011; 58(10):995-1000.
123. Walker S, Levy T, et al. Unitial United Kingdom experience with the use of permanent, biventricular pacemakers. *Europace*, 2000; 2: 233-239.
124. Wells G, Parkash R, Healey JS et al. Cardiac resynchronization therapy: a meta-analysis of randomized controlled trials. *CMAJ* 2011; 183(4):421-9.
125. Whellan DJ, O'Connor CM, et al. Rationale, design, and baseline characteristics of a Program to Assess and Review Trending Information and Evaluate CorRelation to Symptoms in Patients with Heart Failure (PARTNERS HF). *Am Heart J*, November 2008; 156(5): 833-839.
126. Whellan DJ, Ousdigian KT, Al-Khatib SM et al. Combined Heart Failure Device Diagnostics Identify Patients at Higher Risk of Subsequent Heart Failure Hospitalizations Results From PARTNERS HF (Program to Access and Review Trending Information

- and Evaluate Correlation to Symptoms in Patients With Heart Failure) Study. *Journal of the American College of Cardiology* 2010; 55(17):1803-10.
127. Wilton SB, Leung AA, Ghali WA et al. Outcomes of cardiac resynchronization therapy in patients with versus those without atrial fibrillation: a systematic review and meta-analysis. *Heart Rhythm* 2011; 8(7):1088-94.
  128. Woods B, Hawkins N, Mealing S, et al. Individual patient data network meta-analysis of mortality effects of implantable cardiac devices. *Heart*. Nov 2015; 101(22):1800-1806.
  129. Yamokoski LM, Haas GJ, Gans B, and Abraham WT. OptiVol fluid status monitoring with an implantable cardiac device: A heart failure management system. *Expert Rev Med Devices*, November 2007; 4(6): 775-780.
  130. Yancy CW, Jessup M, Bozkurt B, et al. 2013 ACCF/AHA guideline for the management of heart failure: executive summary: a report of the American College of Cardiology Foundation/American heart Association Task Force on practice guidelines. *Circulation*. Oct 15 2013; 128(16): 1810-1852.
  131. Yin J, Hu H, Wang Y, et al. Effects of atrioventricular nodal ablation on permanent atrial fibrillation patients with cardiac resynchronization therapy: a systematic review and meta-analysis. *Clin Cardiol*. Nov 2014; 37(11): 707-715.
  132. Young JB, Abraham WT, Smith AL, et al. Combined cardiac resynchronization and implantable cardioversion defibrillation in advanced chronic heart failure: the MIRACLE ICD Trial. *Jama*. May 28 2003; 289(20):2685-2694.
  133. Ypenburg C, Bax JJ van der Wall EE, et al. Intrathoracic impedance monitoring to predict decompensated heart failure. *Am J Cardiol*, February 2007; 99(4): 554-557.
  134. Yu CM, Abraham WT, Bax J et al. Predictors of response to cardiac resynchronization therapy (PROSPECT)--study design. *Am Heart J* 2005; 149(4):600-5.
  135. Yu CM, Chan JY, Zhang Q et al. Biventricular pacing in patients with bradycardia and normal ejection fraction. *N Engl J Med*. Nov 26, 2009; 361(22)L 2123-2134.
  136. Yu CM, Fang F, Luo XX, et al. Long term follow up results of the pacing to avoid cardiac enlargement (PACE) trial. *Eur J Heart Fail*. Sep 2014; 16(9): 1016-1025.
  137. Yu CM, Wang L, Stadler R et al. Impedance based prediction of CHF admission precedes symptoms in heart failure patients. *Pacing Clin Electrophysiol* 2004; 1(suppl):S213.
  138. Zhang B, Guo J, Zhang G. Comparison of triple-site ventricular pacing versus conventional cardiac resynchronization therapy in patients with systolic heart failure: A meta-analysis of randomized and observational studies. *J Arrhythmia*. 2018;34:55-64.
  139. Zusterzeel R, Seizman KA, Sanders WE, et al. Cardiac resynchronization therapy in women: us food and drug administration meta-analysis of patient level data. *JAMA Intern Med*. Aug 2014; 174(8): 1340-1348.

## **POLICY HISTORY:**

Adopted for Blue Advantage, March 2005

Available for comment May 12-June 27, 2005

Medical Policy Group, August 2006

Medical Policy Group, August 2009  
Available for comment July 17-August 31, 2009  
Medical Policy Group, October 2009  
Available for comment October 20-December 3, 2009  
Medical Policy Group, April 2011  
Available for comment May 11 – June 27, 2011  
Medical Policy Group, November 2011  
Medical Policy Group, April 2012  
Medical Policy Group, April 2013  
Available for comment May 22 through July 6, 2013  
Medical Policy Group, April 2014  
Medical Policy Group, December 2014  
Medical Policy Group, June 2015  
Available for comment June 4 through July 18, 2015  
Medical Policy Group, June 2016  
Medical Policy Group, December 2016  
Medical Policy Group, June 2017  
Medical Policy Group, June 2018  
Available for comment June 11 through July 25, 2018  
Medical Policy Group, December 2018, 2019 CPT Coding Update  
Medical Policy Group, June 2019  
Medical Policy Group, June 2020: Added CPT codes 33207 and 33217  
Medical Policy Group, May 2021

---

*This medical policy is not an authorization, certification, explanation of benefits, or a contract. Eligibility and benefits are determined on a case-by-case basis according to the terms of the member's plan in effect as of the date services are rendered. All medical policies are based on (i) research of current medical literature and (ii) review of common medical practices in the treatment and diagnosis of disease as of the date hereof. Physicians and other providers are solely responsible for all aspects of medical care and treatment, including the type, quality, and levels of care and treatment*

*This policy is intended to be used for adjudication of claims (including pre-admission certification, pre-determinations, and pre-procedure review) in Blue Cross and Blue Shield's administration of plan contracts.*