



BlueCross BlueShield
of Alabama

Name of Blue Advantage Policy:

Cardiac Hemodynamic Monitoring for the Management of Heart Failure in the Outpatient Setting

Policy #: 441
Category: Medicine

Latest Review Date: June 2021
Policy Grade: B

BACKGROUND:

Blue Advantage medical policy does not conflict with Local Coverage Determinations (LCDs), Local Medical Review Policies (LMRPs) or National Coverage Determinations (NCDs) or with coverage provisions in Medicare manuals, instructions or operational policy letters. In order to be covered by Blue Advantage the service shall be reasonable and necessary under Title XVIII of the Social Security Act, Section 1862(a)(1)(A). The service is considered reasonable and necessary if it is determined that the service is:

1. *Safe and effective;*
2. *Not experimental or investigational*;*
3. *Appropriate, including duration and frequency that is considered appropriate for the service, in terms of whether it is:*
 - *Furnished in accordance with accepted standards of medical practice for the diagnosis or treatment of the patient's condition or to improve the function of a malformed body member;*
 - *Furnished in a setting appropriate to the patient's medical needs and condition;*
 - *Ordered and furnished by qualified personnel;*
 - *One that meets, but does not exceed, the patient's medical need; and*
 - *At least as beneficial as an existing and available medically appropriate alternative.*

Routine costs of qualifying clinical trial services with dates of service on or after September 19, 2000 which meet the requirements of the Clinical Trials NCD are considered reasonable and necessary by Medicare. Providers should bill **Original Medicare for covered services that are related to **clinical trials** that meet Medicare requirements (Refer to Medicare National Coverage Determinations Manual, Chapter 1, Section 310 and Medicare Claims Processing Manual Chapter 32, Sections 69.0-69.11).*

POLICY:

Blue Advantage will treat cardiac hemodynamic monitoring for the management of heart failure utilizing inert gas rebreathing, arterial pressure/Valsalva, and implantable direct pressure monitoring of the pulmonary artery as a non-covered benefit and as investigational in the ambulatory care and outpatient setting.

For coverage criteria for cardiac hemodynamic monitoring for the management of heart failure using thoracic bioimpedance, refer to the NCD for Cardiac Output Monitoring by Thoracic Electrical Bioimpedance (TEB) (20.16).

Blue Advantage does not approve or deny procedures, services, testing, or equipment for our members. Our decisions concern coverage only. The decision of whether or not to have a certain test, treatment or procedure is one made between the physician and his/her patient. Blue Advantage administers benefits based on the members' contract and corporate medical policies. Physicians should always exercise their best medical judgment in providing the care they feel is most appropriate for their patients. Needed care should not be delayed or refused because of a coverage determination.

DESCRIPTION OF PROCEDURE OR SERVICE:

A variety of outpatient cardiac hemodynamic monitoring devices are intended to improve quality of life and reduce morbidity for patients with heart failure by decreasing episodes of acute decompensation. Monitors can identify physiologic changes that precede clinical symptoms and thus allow preventive intervention. These devices operate through a variety of mechanisms, including implantable pressure sensors, thoracic bioimpedance measurement, inert gas rebreathing, and estimation of left ventricular end diastolic pressure by arterial pressure during the Valsalva maneuver.

Chronic Heart Failure

Patients with chronic heart failure are at risk of developing acute decompensated heart failure, often requiring hospital admission. Patients with a history of acute decompensation have the additional risk of future episodes of decompensation, and death. Reasons for the transition from a stable, chronic state to an acute, decompensated state include disease progression, as well as acute events such as coronary ischemia and dysrhythmias. While precipitating factors are frequently not identified, the most common preventable cause is noncompliance with medication and dietary regimens.

Management

Strategies for reducing decompensation, and thus the need for hospitalization, are aimed at early identification of patients at risk for imminent decompensation. Programs for early identification of heart failure are characterized by frequent contact with patients to review signs and symptoms with a healthcare provider, education and adjustment of medications as appropriate. These encounters may occur face-to-face in the office or at home, or via cellular or computed technology.

Precise measurement of cardiac hemodynamics is often employed in the intensive care setting to carefully manage fluid status in acutely decompensated heart failure. Transthoracic echocardiography, transesophageal echocardiography (TEE), and Doppler ultrasound are noninvasive methods for monitoring cardiac output on an intermittent basis for the more stable patient but are not addressed herein. A variety of biomarkers and radiological techniques may be used for dyspnea when the diagnosis of acute decompensated heart failure is uncertain.

The criterion standard for hemodynamic monitoring is pulmonary artery catheters and central venous pressure catheters. However, they are invasive, inaccurate and inconsistent in predicting fluid responsiveness. Several studies have demonstrated that catheters fail to improve outcome in critically ill patients and may be associated with harm. To overcome these limitations, multiple techniques and devices have been developed that use complex imaging technology and computer algorithms to estimate fluid responsiveness, volume status, cardiac output and tissue perfusion. Many are intended to be used in outpatient setting but can be used in the emergency department, intensive care unit, and operating room. Five methods are reviewed here: implantable pressure monitoring devices, non-invasive pulmonary fluid monitoring, thoracic bioimpedance, inert gas rebreathing, and arterial waveform during the Valsalva maneuver. The use of last 3 is not widespread because of several limitations including use of proprietary technology making it difficult to confirm their validity and lack of large randomized controlled trials to evaluate treatment decisions guided by these hemodynamic monitors.

This policy refers only to the use of stand-alone cardiac output measurement devices designed for use in ambulatory care and outpatient settings. The use of cardiac hemodynamic monitors or intrathoracic fluid monitors that are integrated into other implantable cardiac devices, including implantable cardioverter defibrillators, cardiac resynchronization therapy devices, and cardiac pacing devices, is addressed in medical policy # 055 – Biventricular Pacemakers (Cardiac Resynchronization Therapy) for the Treatment of Heart Failure.

KEY POINTS:

The most recent literature review was updated through April 5, 2021.

Summary of Evidence

For individuals who have heart failure in outpatient settings who receive hemodynamic monitoring with an implantable pulmonary artery pressure sensor device, the evidence includes randomized controlled trials (RCTs). Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, morbid events, hospitalizations, and treatment-related morbidity. One implantable pressure monitor, the CardioMEMS device, has U.S. Food and Drug Administration approval. The pivotal CHAMPION RCT reported a statistically significant decrease in heart failure-related hospitalizations in patients implanted with CardioMEMS device compared with usual care. However, the results were potentially biased in favor of the treatment group due to use of additional nurse communication to enhance protocol compliance with the device. The manufacture conducted multiple analyses to address the issue of potential bias from the nurse interventions. These were reviewed favorably by FDA. While these analyses demonstrated consistency of benefit from the CardioMEMS device, all such analyses have

methodologic limitations. Early safety data is suggestive of a higher rate of procedural complications, particularly related to pulmonary artery injury. While the U.S. CardioMEMS post-approval study and CardioMEMS European Monitoring Study for Heart Failure (MEMS-HF) study reported a significant decrease in heart-failure related hospitalizations with few device- or system-related complications at 1 year, the impact of nursing interventions remains unclear. Complete 2-year safety outcomes from the CardioMEMS post-approval study are pending, and the serious adverse event rate in the MEMS-HF trial was 8.9%. Given that the intervention is invasive, intended to be used for a highly prevalent condition, in the light of limited safety data, lack of demonstrable mortality benefit and pending questions related to its benefit for reduction in hospitalization, the net benefit remains uncertain. Concerns may be clarified by the ongoing GUIDE-HF RCT that proposes to enroll 3600 patients. The evidence is insufficient to determine that the technology results in an improvement in net health outcomes.

For individuals who have heart failure in outpatient setting who receive hemodynamic monitoring by thoracic bioimpedance, the evidence includes uncontrolled prospective studies and case series. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, morbid events, hospitalizations, and treatment-related morbidity. There is a lack of RCT evidence evaluating whether use of these technologies improves health outcomes over standard active management of heart failure patients. The case series have reported physiologic measurement-related outcomes and/or associations between monitoring information and heart failure exacerbations, but do not provide definitive evidence on device efficacy. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have heart failure in outpatient settings who receive hemodynamic monitoring with inert gas rebreathing or non-invasive pulmonary fluid monitoring, no studies have been identified on clinical validity or clinical utility. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, morbid events, hospitalizations, and treatment-related morbidity. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have heart failure in outpatient settings who receive hemodynamic monitoring of arterial pressure during the Valsalva maneuver, a single study was identified. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, morbid events, hospitalizations, and treatment-related morbidity. The study assessed the use of LVEDP monitoring and reported an 85% sensitivity and an 80% specificity to detect LVEDP greater than 15 mm Hg. The evidence is insufficient to determine the effects of the technology on health outcomes.

Practice Guidelines and Positions Statements

American College of Cardiology et al

The 2017 joint guidelines by the American College of Cardiology, American Heart Association, and Heart Failure Society of America issued joint guidelines on the management of heart failure that offered no recommendations for use of ambulatory monitoring devices.

National Institute for Health and Clinical Excellence

The updated 2018 guidance from the National Institute for Health and Care Excellence (NICE) on chronic heart failure management did not include outpatient hemodynamic monitoring as a recommendation.

In 2013, NICE issued guidance on the insertion and use of implantable pulmonary artery pressure monitors in chronic heart failure. The recommendations concluded that “Current evidence on the safety and efficacy of the insertion and use of implantable pulmonary artery pressure monitors in chronic heart failure is limited in both quality and quantity.”

Heart Failure Society of America

In 2018, the Heart Failure Society of America Scientific Statements Committee published a white paper consensus statement on remote monitoring of patients with heart failure.

The committee concluded that: "Based on available evidence, routine use of external RPM devices is not recommended. Implanted devices that monitor pulmonary arterial pressure and/or other parameters may be beneficial in selected patients or when used in structured programs, but the value of these devices in routine care requires further study."

U.S. Preventive Services Task Force Recommendations

Not applicable

KEY WORDS:

Thoracic electrical bioimpedance, TEB, impedance cardiography, ICD, cardiac output, CO, thermodilution, inert gas rebreathing, BioZ®, Innocor, VeriCor®, Endosure®, Implantable Direct Pulmonary Artery Pressure, Left Ventricular End Diastolic Pressure, LVEDP, Noninvasive Measurement, CardiMEMS, thoracic bioimpedance, TEBCO®, IQ™, Zoe®, Cheetah NICOM®, PhysioFlow®, Cardiography, ZOLL, MicroCor, uCor, HFAMS, Heart Failure and Arrhythmia Management System

APPROVED BY GOVERNING BODIES:

Noninvasive Left Ventricular End Diastolic Pressure Measurement Devices

In June 2004, the VeriCor® (CVP Diagnostics, Boston, MA), a noninvasive left ventricular end diastolic pressure measurement device, was cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process. The FDA determined that this device was substantially equivalent to existing devices for the following indication:

“The VeriCor is indicated for use in estimating non-invasively, left ventricular end-diastolic pressure (LVEDP). This estimate, when used along with clinical signs and symptoms and other patient test results, including weights on a daily basis, can aid the clinician in the selection of further diagnostic tests in the process of reaching a diagnosis and formulating a therapeutic plan when abnormalities of intravascular volume are suspected. The device has been clinically validated in males only. Use of the device in females has not been investigated.”

Thoracic Bioimpedance Devices

Multiple thoracic impedance measurement devices that do not require invasive placement have been cleared for marketing by the U.S. Food and Drug Administration (FDA) 510(k) process. FDA determined that this device was substantially equivalent to existing devices for use for peripheral blood flow monitoring. Table 1 includes a representative list of devices, but is not meant to be comprehensive (FDA product code: DSB).

Table 1: Noninvasive Thoracic Impedance Plethysmography Devices

Device	Manufacturer	Year of FDA Clearance
BioZ® Thoracic Impedance Plethysmograph	SonoSite (Bothell, WA)	2009
Zoe® Fluid Status Monitor	Noninvasive Medical Technologies LLC (Las Vegas, NV)	2004
Cheetah Starling SV	Cheetah Medical Inc.	2008
Physioflow® Signal Morphology-based Impedance Cardiography (SM-ICG™)	Vasocom Inc., now Neumedx Inc. (Bristol, PA)	2008
ReDS™ Wearable System	Sensible Medical Innovations (Philadelphia, PA)	2015

FDA: U.S. Food and Drug Administration.

In addition, several manufacturers market thoracic impedance measurement devices that are integrated into implantable devices.

Non-invasive Pulmonary Fluid Monitoring

In May 2018, the ZOLL uCor (MicroCor) Heart Failure and Arrhythmia Management System (HFAMS) was approved by the FDA through the 510(k) process. The device is described as a “wireless system that employs novel radiofrequency technology to monitor pulmonary fluid levels...ZOLL HFAMS continuously records, stores, and transmits patient data, including Thoracic Fluid Index, heart rate, respiration rate, activity, posture, and heart rhythm.”

Inert Gas Rebreathing Devices

In March 2006, the Innocor® (Innovision, Denmark), an inert gas rebreathing device, was cleared for marketing by FDA through the 510(k) process. FDA determined that this device was substantially equivalent to existing inert gas rebreathing devices for use in computing blood flow. FDA product code: BZG.

Implantable Pulmonary Artery Pressure Sensor Devices

In May 2014, the CardioMEMS™ Champion Heart Failure Monitoring System (CardioMEMS, now Abbott) was approved for marketing by FDA through the premarket approval process. This device consists of an implantable pulmonary artery (PA) sensor, which is implanted in the distal PA, a transvenous delivery system, and an electronic sensor that processes signals from the implantable PA sensor and transmits PA pressure measurements to a secure database. The device originally underwent FDA review in 2011, at which point FDA decided that there was no reasonable assurance that the discussed monitoring system would be effective, particularly in certain subpopulations, although it was agreed that this monitoring system was safe for use in the indicated patient population.

Several other devices that monitor cardiac output by measuring pressure changes in the PA or right ventricular outflow tract have been investigated in the research setting but have not received FDA approval. They include the Chronicle® implantable continuous hemodynamic monitoring device (Medtronic, Minneapolis, MN), which includes a sensor implanted in the right ventricular outflow tract, and the ImPressure® device (Remon Medical Technologies, Caesara, Israel), which includes a sensor implanted in the PA.

Note: This evidence review only addresses use of these techniques in ambulatory care and outpatient settings.

BENEFIT APPLICATION:

Coverage is subject to member's specific benefits. Group specific policy will supersede this policy when applicable.

CURRENT CODING:

CPT Codes:

33289	Transcatheter implantation of wireless pulmonary artery pressure sensor for long-term hemodynamic monitoring, including deployment and calibration of the sensor, right heart catheterization, selective pulmonary catheterization, radiological supervision and interpretation, and pulmonary artery angiography, when performed (Effective 01/01/2019)
93264	Remote monitoring of a wireless pulmonary artery pressure sensor for up to 30 days, including at least weekly downloads of pulmonary artery pressure recordings, interpretation(s), trend analysis, and report(s) by a physician or other qualified health care professional (Effective 01/01/2019)

There is a specific CPT code for bioimpedance:

93701	Bioimpedance-derived physiologic cardiovascular analysis
-------	--

There is a specific CPT code for non-invasive pulmonary monitoring:

0607T	Remote monitoring of an external continuous pulmonary fluid monitoring system, including measurement of radiofrequency-derived pulmonary fluid levels, heart rate, respiration rate, activity, posture, and cardiovascular rhythm (e.g., ECG data), transmitted to a remote 24-hour attended surveillance center; set-up and patient education on use of equipment (Effective 07/01/2020)
0608T	analysis of data received and transmission of reports to the physician or other qualified health care professional (Effective 07/01/2020)

Inert gas rebreathing measurement and LVEDP should be reported using the unlisted code 93799.

There is no specific CPT code for implantable direct pressure monitoring of the pulmonary artery. The unlisted code 93799 would be used.

93799	Unlisted cardiovascular service or procedure
-------	--

REFERENCES:

1. Abraham WT and Adamson P. CardioMEMS completes CHAMPION Clinical Trial Study. Study results indicate that the CardioMEMS implantable hemodynamic monitoring system significantly reduces the leading cause of hospitalizations in the U.S. CardioMEMS™, www.cardiomems.com/content.asp?display=news&view=17.
2. Abraham WT, Adamson PB, Bourge RC et al. Wireless pulmonary artery haemodynamic monitoring in chronic heart failure: a randomised controlled trial. *Lancet* 2011; 377(9766):658-66.
3. Abraham WT, Compton S, Haas G et al. Intrathoracic impedance vs daily weight monitoring for predicting worsening heart failure events: results of the Fluid Accumulation Status Trial (FAST). *Congest Heart Fail* 2011; 17(2):51-5.
4. Abraham J, Bharmi R, Jonsson O, et al. Association of Ambulatory Hemodynamic Monitoring of Heart Failure With Clinical Outcomes in a Concurrent Matched Cohort Analysis. *JAMA Cardiol.* Jun 01 2019; 4(6): 556-563.
5. Abraham WT, Stevenson LW, Bourge RC, et al. Sustained efficacy of pulmonary artery pressure to guide adjustment of chronic heart failure therapy: complete follow-up results from the CHAMPION randomised trial. *Lancet.* Jan 30 2016; 387(10017):453-461.

6. Adamson PB, Abraham WT, Bourge RC, et al. Wireless pulmonary artery pressure monitoring guides management to reduce decompensation in heart failure with preserved ejection fraction. *Circ Heart Fail*. Nov 2014;7(6):935-944.
7. Adamson PB, Abraham WT, Stevenson LW, et al. Pulmonary artery pressure-guided heart failure management reduces 30-day readmissions. *Circ Heart Fail*. Jun 2016;9(6).
8. Adamson PB, Abraham WT, Aaron M et al. CHAMPION trial rationale and design: the long-term safety and clinical efficacy of a wireless pulmonary artery pressure monitoring system. *J Card Fail* 2011; 17(1):3-10.
9. Adamson PB, Gold MR, Bennett T et al. Continuous hemodynamic monitoring in patients with mild to moderate heart failure: results of The Reducing Decompensation Events Utilizing Intracardiac Pressures in Patients with Chronic Heart Failure (REDUCEhf) trial. *Congest Heart Fail* 2011; 17(5):248-54.
10. Amir O, Azzam ZS, Gaspar T, et al. Validation of remote dielectric sensing (ReDS) technology for quantification of lung fluid status: Comparison to high resolution chest computed tomography in patients with and without acute heart failure. *Int J Cardiol*. Oct 15 2016; 221:841-846.
11. Amir O, Ben-Gal T, Weinstein JM, et al. Evaluation of remote dielectric sensing (ReDS) technology-guided therapy for decreasing heart failure re-hospitalizations. *Int J Cardiol*. Aug 1 2017;240:279-284.
12. Anand IS, Greenberg BH, Fogoros RN et al. Design of the Multi-Sensor Monitoring in Congestive Heart Failure (MUSIC) study: prospective trial to assess the utility of continuous wireless physiologic monitoring in heart failure. *Journal of cardiac failure* 2011; 17(1):11-6.
13. Anand IS, Tang WH, Greenberg BH et al. Design and performance of a multisensor heart failure monitoring algorithm: results from the multisensor monitoring in congestive heart failure (MUSIC) study. *Journal of cardiac failure* 2012; 18(4):289-95.
14. Angermann CE, Assmus B, Anker SD, et al. Pulmonary artery pressure-guided therapy in ambulatory patients with symptomatic heart failure: the CardioMEMS European Monitoring Study for Heart Failure (MEMS-HF). *Eur J Heart Fail*. Oct 2020; 22(10): 1891-1901.
15. Belardinelli, R. et al. Comparison of impedance cardiography with thermodilution and direct Fick methods for noninvasive measurements of stroke volume and cardiac output during incremental exercise in patients with ischemic cardiomyopathy, *American Journal of Cardiology*, 1996, Vol. 77, pp. 1293-1301.
16. Bourge RC, Abraham WT, Adamson PB et al. Randomized controlled trial of an implantable continuous hemodynamic monitor in patients with advanced heart failure: the COMPASS-HF study. *Journal of the American College of Cardiology* 2008; 51(11):1073-9.
17. Burns DJP, Arora J, Okunade O, et al. International Consortium for Health Outcomes Measurement (ICHOM): Standardized Patient-Centered Outcomes Measurement Set for Heart Failure Patients. *JACC Heart Fail*. Mar 2020; 8(3): 212-222.

18. CardioMEMSChampion™ Heart Failure Monitoring System: Presentation - CardioMEMS: Oct. 9, 2013. 2013;
<https://wayback.archiveit.org/7993/20170111163201/http://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/MedicalDevices/MedicalDevicesAdvisoryCommittee/CirculatorySystemDevicesPanel/UCM370951.pdf>. Accessed April 17, 2018.
19. CardioMEMS Champion™ HF Monitoring System: FDA Review of P100045/A004FDA Presentation - CardioMEMS: Oct. 9, 2013. 2013;
<https://wayback.archiveit.org/7993/20170111163259/http://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/MedicalDevices/MedicalDevicesAdvisoryCommittee/CirculatorySystemDevicesPanel/UCM370955.pdf>. Accessed April 17, 2018.
20. Chaney, J. et al. Minimally invasive hemodynamic monitoring for the intensivist: Current and emerging technology, *Critical Care Medicine*, October 2002, Vol. 30, No. 10.
21. Christensen P, et al. Thermodilution versus inert gas rebreathing for estimation of effective pulmonary blood flow, *Critical Care Medicine*, January 2000; 28(1): 51-56.
22. Conraads VM, Tavazzi L, Santini M et al. Sensitivity and positive predictive value of implantable intrathoracic impedance monitoring as a predictor of heart failure hospitalizations: the SENSE-HF trial. *European heart journal* 2011; 32(18):2266-73.
23. Damgaard M and Norsh P. Effects of ventilation on cardiac output determined by inert gas rebreathing, *Clinical Physiology and Functional Imaging*, May 2005; 25(3): 142-147.
24. DeMaria, A. et al. Comparative overview of cardiac output measurement methods: Has impedance cardiography come of age?, *Congestive Heart Failure*, March-April 2000; 6(2): 60-73. (Abstract)
25. DeMaria, A. et al. The COST study: A multicenter trial comparing measurement of cardiac output by thoracic electrical bioimpedance with thermodilution, Presentation at the American College of Cardiology 47th Annual Scientific Session, April 1, 1998. (Abstract)
26. Desai AS, Bhimaraj A, Bharmi R, et al. Ambulatory hemodynamic monitoring reduces heart failure hospitalizations in "real-world" clinical practice. *J Am Coll Cardiol*. May 16 2017;69(19):2357-2365.
27. Dickinson, MM, Allen, LL, Albert, NN, et al. Remote Monitoring of Patients With Heart Failure: A White Paper From the Heart Failure Society of America Scientific Statements Committee.. *J. Card. Fail.*, 2018 Oct 12;24(10).
28. Doering, Lynn, et al. Predictors of between-method differences in cardiac output measurement using thoracic electrical bioimpedance and thermodilution, *Critical Care Medicine*, October 1995, Vol. 23 (10), pp. 1667-1673.
29. Drazner, M. et al. Comparison of impedance cardiography with invasive hemodynamic measurements in patients with heart failure secondary to ischemic or nonischemic cardiomyopathy, *The American Journal of Medicine*, April 2002, Vol. 89, No. 8.

30. Dueck, R. et al. Noninvasive cardiac output monitoring, *Chest*, August 2001, Vol. 120, No. 2.
31. Durkin RJ, et al. Noninvasive estimation of pulmonary vascular resistance by stroke index measurement with an inert gas rebreathing technique, *Chest*, July 1994; 106(1): 59-66.
32. FDA. Summary of Safety and Effectiveness Data (SSED) -- CardioMEMS HF System. 2014. Available online at: www.accessdata.fda.gov/cdrh_docs/pdf10/P100045b.pdf. Last accessed March 19, 2020.
33. FDA. 510(k) Clearances. Available at: <https://www.fda.gov/medical-devices/510k-clearances/may-2018-510k-clearances>.
34. Givertz MM, Stevenson LW, Costanzo MR, et al. Pulmonary Artery Pressure-Guided Management of Patients With Heart Failure and Reduced Ejection Fraction. *J Am Coll Cardiol*. Oct 10 2017;70(15):1875-1886.
35. Hassan M, Wagdy K, Kharabish A, et al. Validation of Noninvasive Measurement of Cardiac Output Using Inert Gas Rebreathing in a Cohort of Patients With Heart Failure and Reduced Ejection Fraction. *Circ Heart Fail*. Mar 2017; 10(3).
36. Heist EK, Herre JM, Binkley PF, et al. Analysis of different device-based intrathoracic impedance vectors for detection of heart failure events (from the Detect Fluid Early from Intrathoracic Impedance Monitoring study). *Am J Cardiol*. Oct 15 2014; 114(8):1249-1256.
37. Heywood JT, Jermyn R, Shavelle D et al. Impact of practice-based management of pulmonary artery pressures in 2000 patients implanted with the CardioMEMS Sensor. *Circulation*. 2017 Apr 18; 135(16): 1509-1517.
38. Hirschl, M. et al. Simultaneous comparison of thoracic bioimpedance and arterial pulse waveform-derived cardiac output with thermodilution measurement, *Critical Care Medicine*, June 2000, Vol., 28, No., 6, pp. 1798-1802.
39. International Consortium for Health Outcomes Measurement, Inc (ICHOM). Heart Failure version 1.1.4. Oct 2017. Accessed March 19, 2020.
40. IOM (Institute of Medicine). 2011. *Clinical Practice Guidelines We Can Trust*. Washington, DC: The National Academies Press.
41. Jessup M, Abraham WT, Casey DE, et al. 2009 Writing Group to Review New Evidence and Update the 2005 Guideline for the Management of Patients with Chronic Heart Failure Writing on Behalf of the 2005 Heart Failure. 2009 Focused update: ACCF/AHA Guidelines for the Diagnosis and Management of Heart Failure in Adults: A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines: Developed in Collaboration with the International Society for Heart and Lung Transplantation. *Circulation* 2009; 119: 1977-2016.
42. Jordan HS, Ionnidis JP and Goudas LC. Thoracic electrical bioimpedance 2002. Tufts-New England Medical Center Agency for Healthcare Research and Quality (AHRQ) Evidence-based Practice Center. Available at www.cms.gov. Accessed July 12, 2010.
43. Kamath SA, Drazner MH, Tasissa G et al. Correlation of impedance cardiography with invasive hemodynamic measurements in patients with advanced heart failure: the

BioImpedance CardioGraphy (BIG) substudy of the Evaluation Study of Congestive Heart Failure and Pulmonary Artery Catheterization Effectiveness (ESCAPE) Trial. American heart journal 2009; 158(2):217-23.

44. Krahnke JS, Abraham WT, Adamson PB, et al. Heart failure and respiratory hospitalizations are reduced in patients with heart failure and chronic obstructive pulmonary disease with the use of an implantable pulmonary artery pressure monitoring device. J Card Fail. Mar 2015; 21(3):240-249.
45. Lang CC, Karlin P, Haythe J et al. Ease of noninvasive measurement of cardiac output coupled with peak VO₂ determination at rest and during exercise in patients with heart failure. The American journal of cardiology 2007; 99(3):404-5.
46. Loh JP, Barbash IM, Waksman R. Overview of the 2011 Food and Drug Administration Circulatory System Devices Panel of the Medical Devices Advisory Committee Meeting on the CardioMEMS Champion Heart Failure Monitoring System. Journal of the American College of Cardiology 2013; 61(15):1571-6.
47. Mant J, Al-Mohammad A, Swain S et al. Management of chronic heart failure in adults: synopsis of the National Institute for Health and clinical excellence guideline. Ann Intern Med 2011; 155(4):252-9.
48. McAlister FA, Stewart S, Ferrua S and McMurray JJ. Multidisciplinary strategies for the management of heart failure patients at high risk for admission: A systematic review of randomized trials. JACC, August 2004; 44(4): 810-819.
49. McIntyre KM, Parisi AF, Brown R, et al. Validation and clinical applications of a noninvasive Valsalva response recorder. J Appl Cardiol 1987; 2(2):137-69.
50. McIntyre KM, Vita JA, Lambrew CT, et al. A noninvasive method of predicting pulmonary-capillary wedge pressure. NEJM, December 1992; 327(24): 1715-1720.
51. Moshkovitz Y, et al. Recent developments in cardiac output determination by bioimpedance: Comparison with invasive cardiac output and potential cardiovascular applications, Current Opinion in Cardiology, May 2004; 19(3): 229-237.
52. National Institute for Health and Care Excellence. Insertion and use of implantable pulmonary artery pressure monitors in chronic heart failure, Interventional procedures guidance [IPG463]. 2013; www.nice.org.uk/guidance/ipg463. Accessed April 24, 2018.
53. National Institute for Health and Care Excellence (NICE). Chronic heart failure in adults: diagnosis and management; NICE guideline NG106. Sep 2018. Accessed Apr 2, 2019.
54. Opasich C, Rapezzi C, Lucci D, et al. Precipitating factors and decision-making processes of short-term worsening heart failure despite “optimal” treatment (from the IN-CHF Registry). Am J Cardiol, August 2001; 88(4): 382-387.
55. Packer M, Abraham WT, Mehra MR et al. Utility of impedance cardiography for the identification of short-term risk of clinical decompensation in stable patients with chronic heart failure. Journal of the American College of Cardiology 2006; 47(11):2245-52.
56. Peyton PJ, et al. Agreement of an inert gas rebreathing device with thermodilution and the direct oxygen Fick method in measurement of pulmonary blood flow, Journal of Molecular Modeling, December 2004; 18(5-6): 373-378.

57. Ponikowski P, Voors AA, Anker SD, et al. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *Rev Esp Cardiol (Engl Ed)*. Dec 2016;69(12):1167.
58. Raaijmakers, Esther. A meta-analysis of three decades of validating thoracic impedance cardiography, *Critical Care Medicine*, June 1999, Vol. 27, No. 6.
59. Sharma GVRK, Woods PA, Lambrew CT, et al. Evaluation of a noninvasive system for determining left ventricular filling pressure. *Arch Intern Med* 2002; 162: 2084-2088.
60. Shavelle DM, Desai AS, Abraham WT, et al. Lower Rates of Heart Failure and All-Cause Hospitalizations During Pulmonary Artery Pressure-Guided Therapy for Ambulatory Heart Failure: One-Year Outcomes From the CardioMEMS Post-Approval Study. *Circ Heart Fail*. Aug 2020; 13(8): e006863.
61. Shoemaker, W.C., et al. Multicenter trial of a new thoracic electrical bioimpedance device for cardiac output estimation, *Critical Care Medicine*, December 1994; 22(12): 1907-12.
62. Silber HA, Trost JC, Johnston PV et al. Finger photoplethysmography during the Valsalva maneuver reflects left ventricular filling pressure. *Am J Physiol Heart Circ Physiol* 2012; 302(10):H2043-7.
63. Stevenson LW, Zile M, Bennett TD et al. Chronic ambulatory intracardiac pressures and future heart failure events. *Circ Heart Fail* 2010; 3(5):580-7.
64. Vaduganathan M, DeFilippis EM, Fonarow GC, et al. Postmarketing Adverse Events Related to the CardioMEMS HF System. *JAMA Cardiol*. Nov 1 2017;2(11):1277-1279.
65. Yancy CW, Jessup M, Bozkurt B et al. 2013 ACCF/AHA Guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Circulation* 2013.
66. Yancy CW, Jessup M, Bozkurt B, et al. 2017 ACC/AHA/HFSA Focused Update of the 2013 ACCF/AHA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *J Am Coll Cardiol*. Aug 8 2017;70(6):776-803.
67. Zannad F, Garcia AA, Anker SD, et al. Clinical outcome endpoints in heart failure trials: a European Society of Cardiology Heart Failure Association consensus document. *Eur J Heart Fail*. Oct 2013; 15(10): 1082-94.
68. ZOLL. ZOLL introduces new technology to improve the management of acute heart failure patients. Available at: <https://www.zoll.com/news-releases/2019/06/10/zoll-new-tech-acute-heart-failure-patients-management>.

POLICY HISTORY:

Adopted for Blue Advantage, August 2010

Available for comment August 22 through October 6, 2011

Medical Policy Group, August 2011

Medical Policy Group, July 2012

Medical Policy Group, July 2013
Medical Policy Group, July 2014
Medical Policy Group, September 2014
Medical Policy Group, November 2014
Medical Policy Group, November 2015
Medical Policy Group, May 2016
Medical Policy Group, June 2016
Medical Policy Group, June 2017
Medical Policy Group, May 2018
Medical Policy Group, December 2018: 2019 CPT Coding Update
Medical Policy Group, May 2019
Medical Policy Group, June 2020: Added CPT codes 0607T and 0608T.
Medical Policy Group, June 2021

This medical policy is not an authorization, certification, explanation of benefits, or a contract. Eligibility and benefits are determined on a case-by-case basis according to the terms of the member's plan in effect as of the date services are rendered. All medical policies are based on (i) research of current medical literature and (ii) review of common medical practices in the treatment and diagnosis of disease as of the date hereof. Physicians and other providers are solely responsible for all aspects of medical care and treatment, including the type, quality, and levels of care and treatment.

This policy is intended to be used for adjudication of claims (including pre-admission certification, pre-determinations, and pre-procedure review) in Blue Cross and Blue Shield's administration of plan contracts.