



BlueCross BlueShield  
of Alabama

---

**Name of Blue Advantage Policy:**  
**Hip Resurfacing**

Policy #: 153

Latest Review Date: April 2023

Category: Surgery

---

**BACKGROUND:**

*Blue Advantage medical policy does not conflict with Local Coverage Determinations (LCDs), Local Medical Review Policies (LMRPs) or National Coverage Determinations (NCDs) or with coverage provisions in Medicare manuals, instructions or operational policy letters. In order to be covered by Blue Advantage the service shall be reasonable and necessary under Title XVIII of the Social Security Act, Section 1862(a)(1)(A). The service is considered reasonable and necessary if it is determined that the service is:*

1. *Safe and effective;*
2. *Not experimental or investigational\*;*
3. *Appropriate, including duration and frequency that is considered appropriate for the service, in terms of whether it is:*
  - *Furnished in accordance with accepted standards of medical practice for the diagnosis or treatment of the patient's condition or to improve the function of a malformed body member;*
  - *Furnished in a setting appropriate to the patient's medical needs and condition;*
  - *Ordered and furnished by qualified personnel;*
  - *One that meets, but does not exceed, the patient's medical need; and*
  - *At least as beneficial as an existing and available medically appropriate alternative.*

*\*Routine costs of qualifying clinical trial services with dates of service on or after September 19, 2000 which meet the requirements of the Clinical Trials NCD are considered reasonable and necessary by Medicare. Providers should bill **Original Medicare** for covered services that are related to **clinical trials** that meet Medicare requirements (Refer to Medicare National Coverage Determinations Manual, Chapter 1, Section 310 and Medicare Claims Processing Manual Chapter 32, Sections 69.0-69.11).*

## **POLICY:**

**Blue Advantage** will treat **metal-on-metal total hip resurfacing with an FDA-approved device system** as a **covered** benefit as an alternative to total hip replacement when the patient:

- Is a candidate for total hip replacement; AND
- Is likely to outlive a traditional prosthesis; AND
- Does not have a contraindication for total hip resurfacing (See below).

Contraindications for total hip resurfacing:

- Bone stock inadequate to support the device due to:
  - Severe osteopenia or a family history of severe osteoporosis or severe osteopenia
  - Osteonecrosis or avascular necrosis with more than 50% involvement of the femoral head
  - Multiple cysts of the femoral head (more than 1 cm)
- Skeletal immaturity
- Vascular insufficiency, muscular atrophy, or neuromuscular disease severe enough to compromise implant stability or postoperative recovery
- Known moderate to severe renal insufficiency
- Severely overweight
- Known or suspected metal sensitivity
- Immunosuppressed or receiving high doses of corticosteroids
- Females of child bearing age due to unknown effects of the fetus of metal ion release

**Blue Advantage** will treat **partial hip resurfacing with an FDA-approved device** as a **covered** benefit in patients with osteonecrosis of the femoral head who have one or more contraindications for metal-on-metal implants and meet the following criteria:

- The patient is a candidate for total hip replacement; and
- Is likely to outlive a traditional prosthesis; and
- The patient has known or suspected metal sensitivity or concern about potential effects of metal ions; and
- There is no more than 50% involvement of the femoral head; and
- There is minimal change in acetabular cartilage or articular cartilage space identified on radiography.

**Blue Advantage** will treat **all other types and applications** of hip resurfacing as a **non-covered** benefit and as **investigational**.

*Blue Advantage does not approve or deny procedures, services, testing, or equipment for our members. Our decisions concern coverage only. The decision of whether or not to have a certain test, treatment or procedure is one made between the physician and his/her patient. Blue Advantage administers benefits based on the members' contract and medical policies. Physicians should always exercise their best medical judgment in providing the care they feel is most appropriate for their patients. Needed care should not be delayed or refused because of a coverage determination.*

## **DESCRIPTION OF PROCEDURE OR SERVICE:**

Hip resurfacing is an alternative to total hip arthroplasty (THA, also known as hip replacement) for patients with advanced arthritis of the hip. Total hip resurfacing (THR) describes the placement of a shell that covers the femoral head together with implantation of an acetabular cup in patients with painful hip joints. Partial hip resurfacing is considered a treatment option for avascular necrosis with collapse of the femoral head. Available prostheses are metal-on-metal devices.

THR has been investigated in patients with osteoarthritis, rheumatoid arthritis, and advanced avascular necrosis as an alternative to THA, particularly in young active patients who would potentially outlive a total hip prosthesis. Therefore, hip resurfacing could be viewed as a time-buying procedure to delay the need for a THA. Proposed advantages of THR compared with THA include preservation of the femoral neck and femoral canal, thus facilitating revision or conversion to a THR, if required. In addition, the resurfaced head is more similar in size to the normal femoral head, thus increasing the stability and decreasing the risk of dislocation compared with THA.

THR has undergone various evolutions, with modifications in prosthetic design and composition and implantation techniques. For example, similar to total hip prostheses, the acetabular components of THR have been composed of polyethylene. However, over time it became apparent that device failure was frequently related to the inflammatory osteolytic reaction to polyethylene debris wear particles. Metal acetabular components have since been designed to improve implant longevity. Sensitivity to wear particles from metal-on-metal chromium and cobalt implant components are of increasing concern.

## **KEY POINTS:**

The most recent literature update was performed through February 16, 2023.

### **Summary of Evidence**

For individuals who have an indication for hip replacement who would outlive a traditional prosthesis and have no contraindication for hip resurfacing who receive a metal-on-metal total hip resurfacing device, the evidence includes RCTs, numerous large observational studies, large registry studies, and systematic reviews. Relevant outcomes are symptoms, change in disease status, functional outcomes, health status measures, quality of life, and treatment-related morbidity. The efficacy of total hip resurfacing performed with current techniques is similar to that for THA over the short-to-medium term, and total hip resurfacing may permit easier conversion to a THA for younger patients expected to outlive their prosthesis. Based on potential ease of revision of total hip resurfacing compared with THA, current evidence supports conclusions that hip resurfacing presents a reasonable alternative for active patients who are considered too young for THA when performed by surgeons experienced in the technique. The literature on adverse events (e.g., metallosis, pseudotumor formation, implant failure) is evolving as longer follow-up data become available. Due to the uncertain risk with metal-on-metal implants, the risk-benefit ratio needs to be considered carefully on an individual basis. In addition, emerging evidence has suggested an increased risk of failure in women, possibly due to smaller implant size. Therefore, these factors should also be considered in the overall patient

evaluation for total hip resurfacing, and patients should make an informed choice with their treating physicians. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have an indication for hip replacement who would outlive a traditional prosthesis and have no contraindication for hip resurfacing who receive a partial hip resurfacing device, the evidence includes a comparative study. Relevant outcomes are symptoms, change in disease status, functional outcomes, health status measures, QOL, and treatment-related morbidity. Although evidence has shown better outcomes with THR than with partial hip resurfacing, partial hip resurfacing would be appropriate in younger patients with osteonecrosis who have contraindications for a metal-on-metal prosthesis. These factors should be considered in the overall patient evaluation for THR, and patients should make an informed choice with their treating physicians. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

## **Practice Guidelines and Position Statements**

### **Hip Society**

The Hip Society (2012) published an algorithmic approach to the diagnosis and management of metal-on-metal (MoM) arthroplasty (total hip arthroplasty [THA], total hip resurfacing [THR]. The review indicated that adverse local tissue reactions to metal debris are escalating and that all arthroplasty patients returning for follow-up should be queried for pain, discomfort, or compromise of function. Symptomatic patients should be evaluated for all intra-articular and extra-articular causes of pain, including aseptic loosening, sepsis, component malposition, or fluid collections and/or masses about the hip. The Hip Society stated that there is still a role for MoM resurfacing arthroplasty in select patients groups. The ideal candidate is a man younger than age 55 with osteoarthritis and a femoral head size larger than 50 mm. Another relative indication is the need or desire to return to a very high activity level at work or in recreation. Contraindications to MoM resurfacing include known or suspected metal sensitivity; moderate or worse renal function; women who may become pregnant; osteoporosis; large cysts; and avascular necrosis more than 50%.

### **American Academy of Orthopaedic Surgeons**

In 2010, the American Academy of Orthopaedic Surgeons published a technology overview on MoM hip resurfacing. To compare revision rates between MoM hip resurfacing and THA, the Academy analyzed three joint registries, which indicated that patients who received THR were at greater risk for revision than patients who received THA. One registry suggested that younger men may have a lower revision rate after THR than THA, although the available data were not found to clearly establish an advantage for this subgroup. There was no conclusive evidence on predictors of successful or unsuccessful outcomes.

### **National Institute for Health and Care Excellence**

The NICE (2014) updated its guidance on THA and THR for end-stage arthritis of the hip. The NICE concluded that both THA and THR were options for treating end-stage arthritis of the hip, although clinicians may be more likely to offer resurfacing arthroplasty to men than to women because of higher revision rates observed in women. The NICE concluded that THA was more effective and less costly than THR in all analyses, that the revision rate was the most important

key driver of costs and quality-adjusted life years, and that because the predicted revision rate of THA was less than 5% at ten years in the population for whom both THA and THR were suitable, the revision rate standard for THR should be the same as that for THA. The NICE recommended specific prostheses for THA and THR only if the prostheses have revision rates of 5% or less at ten years.

#### **U.S. Preventive Services Task Force Recommendations**

Not applicable.

#### **KEY WORDS:**

Total hip resurfacing, partial hip resurfacing, total hip arthroplasty, Buechel-Pappas Integrated Total Hip Replacement, Conserve Plus, Cormet 2000, Birmingham hip resurfacing device, BHR

#### **APPROVED BY GOVERNING BODIES:**

In 2006, FDA approved the Birmingham Hip Resurfacing (BHR; Smith & Nephew Orthopaedics, Cordova, TN) system, a MoM resurfacing system, through the premarket approval (PMA) process for use in patients requiring primary hip resurfacing arthroplasty for non-inflammatory or inflammatory arthritis. This decision was primarily based on a series of 2203 patients (2385 hips) who received this device by a single surgeon in England. A number of post-approval requirements were agreed to, including the following items:

- Study longer-term safety and effectiveness through 10-year follow-up of the first consecutive 350 cases in the 2385 hip case cohort that was part of the PMA.
- Study the “learning curve” and the longer-term safety and effectiveness of the BHR in the United States by studying 350 patients at up to 8 sites where clinical and radiographic data will be assessed annually through 5 years and at 10 years. Also, determine cobalt and chromium serum concentration and renal function in these patients at 1, 4, and 10 years.
- Implement a training program to provide clinical updates to investigators.

Two additional MoM hip resurfacing systems have been approved: in 2007, the Cormet™ Hip Resurfacing System (Corin, Tampa, FL) and, in 2009, the Conserve® Plus Total Hip Resurfacing System (MicroPort Orthopedics, Arlington, TN). Both implants were approved for skeletally mature patients with either non-inflammatory degenerative arthritis (e.g., osteoarthritis and avascular necrosis) or inflammatory arthritis (e.g., rheumatoid arthritis). (Note: patients with the latter arthritis might be individuals who, due to younger age or increased activity level, may not be suitable for traditional THA because it would increase the possibility of requiring ipsilateral hip joint revision.)

Various devices have been cleared for marketing by FDA through the 510(k) process for partial hip (femoral) resurfacing. Some surgeons may be using a femoral resurfacing component together with an acetabular cup (total arthroplasty component) as an "off-label" application.

**BENEFIT APPLICATION:**

Coverage is subject to member's specific benefits. Group specific policy will supersede this policy when applicable.

**CURRENT CODING:**

There is no specific CPT code for total hip resurfacing. The American Academy of Orthopaedic Surgeon's coding committee has written several articles stating that this procedure should be reported with the regular total hip CPT code 27130.

**CPT:**

27299	Unlisted procedure, pelvis or hip joint
27130	Arthroplasty, acetabular and proximal femoral prosthetic replacement [total hip replacement], with or without autograft or allograft

**HCPCS:**

S2118	Metal-on-metal total hip resurfacing, including acetabular and femoral components
-------	---

**REFERENCES:**

1. Amstutz HC, Le Duff MJ, Campbell PA et al. Clinical and radiographic results of metal-on-metal hip resurfacing with a minimum ten-year follow-up. J Bone Joint Surg Am 2010; 92(16):2663-2671.
2. Australian Orthopedic Association. National Joint Replacement Registry Annual Report. 2006.
3. Azam MQ, McMahon S, Hawdon G, et al. Survivorship and clinical outcome of Birmingham hip resurfacing: a minimum ten years' follow-up. Int Orthop. Jan 2016; 40(1):1-7.
4. Ball ST, LeDuff MJ and Amstutz HC. Early results of conversion of a failed femoral component in hip resurfacing arthroplasty. J Bone Joint Surg Am, April 2007; 89(4): 735-741.
5. Beaulé PE, Amstutz HC, Le Duff M et al. Surface arthroplasty for osteonecrosis of the hip: hemiresurfacing versus metal-on-metal hybrid resurfacing. J Arthroplasty 2004; 19(8 Suppl 3):54-58.
6. Bourget-Murray J, Watt Kearns SJ, Piroozfar S, et al. Birmingham Hip Resurfacing for osteoarthritis - a Canadian retrospective cohort study with a minimum 10-year follow-up. Can J Surg. 2022; 65(3): E296-E302.
7. Daniel J, Pradhan C, Ziaee H, et al. Results of Birmingham hip resurfacing at 12 to 15 years: a single-surgeon series. Bone Joint J. Oct 2014; 96- B(10):1298-1306.
8. de Steiger RN, Miller LN, Prosser GH et al. Poor outcome of revised resurfacing hip arthroplasty. Acta Orthop 2010; 81(1):72-76.

9. Food and Drug Administration. P040033: Birmingham Hip Resurfacing (BHR) System. 2006; [http://www.accessdata.fda.gov/cdrh\\_docs/pdf4/p040033a.pdf](http://www.accessdata.fda.gov/cdrh_docs/pdf4/p040033a.pdf).
10. Garbuz DS, Tanzer M, Greidanus NV et al. The John Charnley award: metal-on-metal hip resurfacing versus large-diameter head metal-on-metal total hip arthroplasty: a randomized clinical trial. Clin Orthop Relat Res Feb 2010; 468(2):318-325.
11. Grcula MJ. Resurfacing arthroplasty in osteonecrosis of the hip. Orthop Clin North Am 2005; 36(2):231-242, x.
12. Gross TP, Liu F, Webb LA. Clinical outcome of the metal-on-metal hybrid Corin Cormet 2000 hip resurfacing system: an up to 11-year follow-up study. J Arthroplasty 2012; 27(4):533-538 e531.
13. Haddad FS, Konan S, Tahmassebi J. A prospective comparative study of cementless total hip arthroplasty and hip resurfacing in patients under the age of 55 years: a ten-year follow-up. Bone Joint J. May 2015;97-B(5):617-622.
14. IOM (Institute of Medicine). 2011. Clinical Practice Guidelines We Can Trust. Washington, DC: The National Academies Press.
15. Jiang Y, Zhang K, Die J et al. A systematic review of modern metal-on-metal total hip resurfacing vs standard total hip arthroplasty in active young patients. J Arthroplasty 2011; 26(3):419-426.
16. Kim PR, Beaulé PE, Lafamme Gy et al. Causes of early failure in a multicenter clinical trial of hip resurfacing. J Arthroplasty 2008; 23(6 Suppl 1):44-49.
17. Kwon YM, Ostlere SJ, McLardy-Smith P et al. Asymptomatic pseudotumors after metal-on-metal hip resurfacing arthroplasty: prevalence and metal ion study. J Arthroplasty 2011; 26(4):511-518.
18. Kumar P, Ksheersagar V, Aggarwal S, et al. Complications and mid to long term outcomes for hip resurfacing versus total hip replacement: a systematic review and meta-analysis. Eur J Orthop Surg Traumatol. Aug 25 2022.
19. Lavigne M, Therrien M, Nantel J et al. The John Charnley award: the functional outcome of hip resurfacing and large-head THA is the same: a randomized, double-blind study. Clin Orthop Relat Res. Feb 2010; 468(2): 326-336.
20. Lass R, Bechler U, Springer B, et al. Midterm results of the Birmingham hip resurfacing: a single-surgeon series. Arch Orthop Trauma Surg. Feb 2023; 143(2): 1041-1048.
21. Lombardi AV, Jr., Barrack RL, Berend KR, et al. The Hip Society: algorithmic approach to diagnosis and management of metal-on-metal arthroplasty. J Bone Joint Surg Br. Nov 2012; 94(11 Suppl A):14-18.
22. Marker DR, Strimbu K, McGrath MS et al. Resurfacing versus conventional total hip arthroplasty-review of comparative clinical and basic science studies. Bull NYU Hosp Jt Dis 2009; 67(2):120-127.
23. Matharu GS, McBryde CW, Pynsent WB, et al. The outcome of the Birmingham Hip Resurfacing in patients aged < 50 years up to 14 years post-operatively. Bone Joint J. Sep 2013; 95-B(9):1172-1177.
24. McGrath MS, Marker DR, Seyler TM et al. Surface replacement is comparable to primary total hip arthroplasty. Clin Orthop Related Res. Jan 2009; 467(1):94-100.

25. McGrory B, Barrack R, Lachiewicz PF, et al. Modern metal-on-metal hip resurfacing. *J Am Acad Orthop Surg*. May 2010;18(5):306-314.
26. Mont MA, Seyler TM, Ragland PS, Starr R, et al. Gait analysis of patients with resurfacing hip arthroplasty compared with hip osteoarthritis and standard total hip arthroplasty. *J Arthroplasty*, January 2007; 22(1): 100-108.
27. Mont MA, Seyler TM, Ulrich SD, et al. Effect of changing indications and techniques on total hip resurfacing. *Clin Orthop Relat Res*, December 2007; 465: 63-70.
28. Murray DW, Grammatopoulos G, Pandit H, et al. The ten-year survival of the Birmingham hip resurfacing: an independent series. *J Bone Joint Surg Br*. Sep 2012; 94(9):1180-1186.
29. National Institute for Health and Care Excellence (NICE). Total hip replacement and resurfacing arthroplasty for end-stage arthritis of the hip [TA304]. 2014; [www.nice.org.uk/guidance/ta304](http://www.nice.org.uk/guidance/ta304).
30. Nunley RM, De la Valle CJ, Barrack RL. Is patient selection important for hip resurfacing? *Clin Orthop Relat Res* 2009; 467(1):56-65.
31. Nunley RM, Zhu J, Brooks PJ et al. The learning curve for adopting hip resurfacing among hip specialists. *Clin Orthop Relat Res* Feb 2010; 468(2):382-391.
32. Ollivere B, Darrah C, Barker T et al. Early clinical failure of the Birmingham metal-on-metal hip resurfacing is associated with metallosis and soft-tissue necrosis. *J Bone Joint Surg Br* 2009; 91(8):1025-1030.
33. Pailhe R, Matharu GS, Sharma A, et al. Survival and functional outcome of the Birmingham Hip Resurfacing system in patients aged 65 and older at up to ten years of follow-up. *Int Orthop*. Jun 2014; 38(6):1139-1145.
34. Quesada MJ, Marker DR, Mont MA. Metal-on-metal hip resurfacing: advantages and disadvantages. *J Arthroplasty* 2008; 23(7 Suppl); 69-73.
35. Reito A, Puolakka T, Elo P, et al. Outcome of Birmingham hip resurfacing at ten years: role of routine whole blood metal ion measurements in screening for pseudotumours. *Int Orthop*. Nov 2014; 38(11):2251-2257.
36. Steffen RT, Pandit HP, Palan J et al. The five-year results of the Birmingham Hip Resurfacing arthroplasty: an independent series. *J Bone Joint Surg Br* 2008; 90(4):436-441.
37. Stoney J, Graves SE, de Steiger RN, et al. Is the Survivorship of Birmingham Hip Resurfacing Better Than Selected Conventional Hip Arthroplasties in Men Younger Than 65 Years of Age? A Study from the Australian Orthopaedic Association National Joint Replacement Registry. *Clin Orthop Relat Res*. Nov 2020; 478(11): 2625-2636.
38. Stulberg BN, Fitts SM, Zadzilka JD et al. Resurfacing arthroplasty for patients with osteonecrosis. *Bull NYU Hosp Jt Dis* 2009; 67(2):138-141.
39. Su EP, Ho H, Bhal V, et al. Results of the First U.S. FDA-Approved Hip Resurfacing Device at 10-Year Follow-up. *J Bone Joint Surg Am*. Jul 21 2021; 103(14): 1303-1311.
40. Van Der Straeten C, Gross TP, Amstutz H, et al. Hip resurfacing arthroplasty in young patients: international high-volume centres' report on the outcome of 11,382 metal-on-metal hip resurfacing arthroplasties in patients ≤50 years at surgery. *Hip Int*. May 2022; 32(3): 353-362.



41. Vendittoli PA, Lavigne M, Roy AG, et al. A prospective randomized clinical trial comparing metal-on-metal total hip arthroplasty and metal-on-metal total hip resurfacing in patients less than 65 years old. *Hip Int.* 2006; 16 Suppl 4:73-81.
42. Williams DH, Greidanus NV, Masri BA et al. Prevalence of pseudotumor in asymptomatic patients after metal-on-metal hip arthroplasty. *J Bone Joint Surg Am* 2011; 93(23):2164-2171.

## **POLICY HISTORY:**

Adopted for Blue Advantage, March 2005

Available for comment May 1-June 14, 2005

Medical Policy Group, March 2006

Medical Policy Group, January 2007

Available for comment February 6-March 22, 2007

Medical Policy Group, January 2009

Medical Policy Group, December 2009

Medical Policy Group, January 2010

Available for comment February 6-March 22, 2010

Medical Policy Group, October 2012

Medical Policy Group, November 2013

Medical Policy Group, September 2014

Medical Policy Group, October 2015

Medical Policy Group, September 2017

Medical Policy Group, May 2018

Medical Policy Group, May 2019

Medical Policy Group, May 2020

Medical Policy Group, April 2021

Medical Policy Group, April 2022

Medical Policy Group, April 2023

---

*This medical policy is not an authorization, certification, explanation of benefits, or a contract. Eligibility and benefits are determined on a case-by-case basis according to the terms of the member's plan in effect as of the date services are rendered. All medical policies are based on (i) research of current medical literature and (ii) review of common medical practices in the treatment and diagnosis of disease as of the date hereof. Physicians and other providers are solely responsible for all aspects of medical care and treatment, including the type, quality, and levels of care and treatment.*

*This policy is intended to be used for adjudication of claims (including pre-admission certification, pre-determinations, and pre-procedure review) in Blue Cross and Blue Shield's administration of plan contracts.*