



**BlueCross BlueShield
of Alabama**

Name of Blue Advantage Policy:

Phototherapy for the Treatment of Skin Disorders

Policy #: 301

Latest Review Date: January 2025

Category: Medical/DME

BACKGROUND:

Blue Advantage medical policy does not conflict with Local Coverage Determinations (LCDs), Local Medical Review Policies (LMRPs) or National Coverage Determinations (NCDs) or with coverage provisions in Medicare manuals, instructions or operational policy letters. In order to be covered by Blue Advantage the service shall be reasonable and necessary under Title XVIII of the Social Security Act, Section 1862(a)(1)(A). The service is considered reasonable and necessary if it is determined that the service is:

1. *Safe and effective;*
2. *Not experimental or investigational*;*
3. *Appropriate, including duration and frequency that is considered appropriate for the service, in terms of whether it is:*
 - *Furnished in accordance with accepted standards of medical practice for the diagnosis or treatment of the patient's condition or to improve the function of a malformed body member;*
 - *Furnished in a setting appropriate to the patient's medical needs and condition;*
 - *Ordered and furnished by qualified personnel;*
 - *One that meets, but does not exceed, the patient's medical need; and*
 - *At least as beneficial as an existing and available medically appropriate alternative.*

Routine costs of qualifying clinical trial services with dates of service on or after September 19, 2000 which meet the requirements of the Clinical Trials NCD are considered reasonable and necessary by Medicare. Providers should bill **Original Medicare for covered services that are related to **clinical trials** that meet Medicare requirements (Refer to Medicare National Coverage Determinations Manual, Chapter 1, Section 310 and Medicare Claims Processing Manual Chapter 32, Sections 69.0-69.11).*

POLICY:

Blue Advantage will treat **ultraviolet A or B therapy (CPT 96900)** as a **covered** benefit in the treatment of the following conditions:

- Chronic urticaria
- Eczema (atopic dermatitis)
- Lichen planus
- Mycosis fungoides (cutaneous T-cell lymphoma)
- Pityriasis lichenoides
- Pityriasis rosea
- Pruritus of renal failure
- Vitiligo
- Localized scleroderma

Blue Advantage will treat **ultraviolet B with the addition of topical coal tar** (also known as Goeckerman treatment) **or petrolatum (CPT 96910)** as a **covered benefit** for **severe psoriasis** (defined as psoriasis that affects more than 10% of the body surface area).

Blue Advantage will treat **ultraviolet B with the addition of topical coal tar** (also known as Goeckerman treatment) **or petrolatum** as a **non-covered benefit** and as **investigational for all other indications**.

Blue Advantage will treat **ultraviolet B light therapy (CPT 96900) administered in the home** as a **covered** benefit for the following conditions and when conducted under a physician's supervision with regularly scheduled exams:

- Atopic dermatitis-mild to moderate forms when standard treatment has failed
- Lichen planus
- Mycosis fungoides
- Pityriasis lichenoides
- Pruritus of hepatitis disease
- Pruritus of renal failure
- Severe atopic dermatitis

Blue Advantage will treat **ultraviolet B light therapy (CPT 96900) administered in the home** as a **non-covered benefit for conditions not listed above**.

Blue Advantage will treat **Psoralens and ultraviolet A light (PUVA) therapy (CPT 96912)** as a **covered** benefit for the following conditions:

- Acrodermatitis continua
- Acute/chronic pityriasis lichenoides
- Eczema (atopic dermatitis)
- Lichen planus
- Mycosis fungoides (cutaneous T-cell lymphoma)
- Nummular dermatitis
- Palmoplantar pustulosis

- Parapsoriasis
- Poikiloderma vasculare
- Psoriasis
- Pustulosis palmaris
- Vitiligo

Blue Advantage will treat **photochemotherapy (Goeckerman and/or PUVA) for severe photoresponsive dermatoses requiring at least 4-8 hours of care under direct supervision of the physician (CPT 96913)** as a **covered benefit** for the following conditions:

- Acrodermatitis continua
- Acute/chronic pityriasis lichenoides
- Eczema (atopic dermatitis)
- Lichen planus
- Mycosis fungoides (cutaneous T-cell lymphoma)
- Psoriasis
- Pustulosis palmaris
- Vitiligo

Blue Advantage will treat **excimer laser treatment of vitiligo of the face, neck, trunk, abdomen, back and/or proximal limbs** as a **covered benefit for up to three sessions per week for 12 weeks.**

Blue Advantage will treat **excimer laser treatment of vitiligo of the distal limbs and bony prominences (i.e. fingers, wrists, elbows, knees)** as a **non-covered benefit** and as **investigational.**

Blue Advantage does not approve or deny procedures, services, testing, or equipment for our members. Our decisions concern coverage only. The decision of whether or not to have a certain test, treatment or procedure is one made between the physician and his/her patient. Blue Advantage administers benefits based on the members' contract and medical policies. Physicians should always exercise their best medical judgment in providing the care they feel is most appropriate for their patients. Needed care should not be delayed or refused because of a coverage determination.

DESCRIPTION OF PROCEDURE OR SERVICE:

Phototherapy is defined as the exposure to nonionizing radiation for therapeutic benefit. It may involve exposure to UVA UVB or various combinations of UVA and UVB radiation. UV light therapy, including phototherapy and photochemotherapy, is used for the treatment of certain skin conditions.

Targeted phototherapy may also be used in specific conditions that have not responded to standard therapies describes the use of ultraviolet light that can be focused on specific body areas or lesions. In contrast, photochemotherapy or psoralens in conjunction with UVA is the

therapeutic use of radiation in combination with a photosensitizing chemical. Treatment with these modalities may involve partial or whole-body exposure. Photochemotherapy has been used for a large number of skin diseases but confirmed data of its usefulness is available in only a relatively few. PUVA uses a psoralen derivative in conjunction with long wavelength UVA light (sunlight or artificial) for photochemotherapy of skin conditions.

Targeted phototherapy with handheld lamps or lasers is also being evaluated. Potential advantages of targeted phototherapy include the ability to use higher treatment doses and to limit exposure to surrounding tissue. Original UVB devices consisted of a Phillips TL-01 fluorescent bulb with a maximum wavelength (λ_{max}) at 311 nm. Subsequently, xenon chloride (XeCl) lasers and lamps were developed as targeted UVB treatment devices; these devices generate monochromatic or very narrowband (NB) radiation with a λ_{max} of 308 nm. Targeted phototherapy devices are directed at specific lesions or affected areas, thus limiting exposure to the surrounding normal tissues. They may therefore allow higher dosages compared with a light box, which could result in fewer treatments.

There are numerous medical and surgical treatments aimed at decreasing disease progression and/or attaining repigmentation. Topical corticosteroids, alone or in combination with topical vitamin D3 analogues, are common first-line treatment for vitiligo. Alternative first-line therapies include topical calcineurin inhibitors, systemic steroids, and topical antioxidants. Treatment options for vitiligo recalcitrant to first-line therapy include, among others, light box therapy with narrowband ultraviolet B (NB-UVB) and psoralen plus ultraviolet A (PUVA) and targeted light therapy.

Light therapy for vitiligo includes both targeted phototherapy and photochemotherapy with psoralen plus PUVA. Vitiligo is an idiopathic skin disorder that causes depigmentation of sections of skin, most commonly on the extremities. Depigmentation occurs because melanocytes are no longer able to function properly. The cause of vitiligo is unknown; it is sometimes considered an autoimmune disease. The most common form of the disorder is nonsegmental vitiligo (NSV) in which depigmentation is generalized, bilateral, symmetrical, and increases in size over time. In contrast, segmental vitiligo (SV), also called asymmetric or focal vitiligo, covers a limited area of skin. The typical natural history of vitiligo involves stepwise progression with long periods in which the disease is static and relatively inactive, and relatively shorter periods in which areas of pigment loss increase.

PUVA uses a psoralen derivative in conjunction with long-wavelength UVA light (sunlight or artificial) for photochemotherapy of skin conditions. Psoralens are tricyclic furocoumarin that occur in certain plants and can also be synthesized. They are available in oral and topical forms. Oral PUVA is generally given 1.5 hours before exposure to UVA radiation. Topical PUVA therapy refers the direct application of psoralen to the skin with subsequent exposure to UVA light. With topical PUVA, UVA exposure is generally administered within thirty minutes of psoralen application. No topical psoralen formulation is currently available in the US.

Excimer laser is a form of ultraviolet laser proposed for the treatment of various dermatologic conditions including atopic dermatitis, psoriasis and vitiligo. Laser therapy provides intense UVB light to a limited area of skin, providing the potential benefit of more rapid clinical

response from targeted phototherapy while avoiding the side effects of ultraviolet light on unaffected skin.

KEY POINTS:

The most recent literature update was performed through October 29, 2024.

Summary of Evidence

For individuals who have vitiligo who receive targeted phototherapy, the evidence includes systematic reviews of randomized controlled trials (RCTs), 2 individual RCTs, and 2 retrospective studies. Relevant outcomes are a change in disease status, quality of life (QOL), and treatment-related morbidity. Individual studies tend to have small sample sizes, and few were designed to isolate the effect of laser therapy. Two meta-analyses were attempted; however, results from a meta-analysis could not be verified because the selected studies were not available in English, and 1 estimate was imprecise due to the small number of studies and participants. RCTs have shown targeted phototherapy to be associated with statistically significant improvements in Vitiligo Area Scoring Index (VASI) scores and/or repigmentation compared to alternate treatment options in some studies, but the results are inconsistent. Overall, there is a lack of clinical trial evidence that compares targeted phototherapy with more conservative treatments or no treatment/placebo.

For individuals who have vitiligo who have not responded to conservative therapy who receive PUVA (photochemotherapy), the evidence includes systematic reviews and RCTs. Relevant outcomes are a change in disease status, quality of life, and treatment-related morbidity. There is some evidence from randomized studies, mainly those published before 1985, that PUVA is more effective than a placebo for treating vitiligo. When compared with narrowband UVB (NB-UVB) in meta-analyses, results have shown that individuals receiving NB-UVB experienced higher rates of repigmentation than individuals receiving PUVA, though the differences were not statistically significant. Based on the available evidence and clinical guidelines, PUVA may be considered in individuals with vitiligo who have not responded adequately to conservative therapy.

Practice Guidelines and Position Statements

Vitiligo Working Group

The Vitiligo Working Group is supported by the National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS), part of the National Institutes of Health (NIH). In 2017, the group published guidelines on current and emerging treatments for vitiligo. The Working Group indicated that psoralens with UVA has largely been replaced by NB-UVB, but that “PUVA may be considered in individuals with darker Fitzpatrick skin phototypes or those with treatment-resistant vitiligo (level I evidence).” The Working Group also stated that “Targeted phototherapy (excimer lasers and excimer lamps) can be considered when <10% of body surface area is affected (level II evidence).”

U.S. Preventive Services Task Force Recommendations

Not applicable.

KEY WORDS:

Phototherapy, photochemotherapy, UVA, UVB, PUVA, ultraviolet A, ultraviolet B, excimer laser phototherapy, excimer laser, 308-nm excimer laser, 308-nm xenon chloride excimer laser, vitiligo, psoralen plus, atopic dermatitis, Handisol II[®], XTRAC[®], Xenon monochloride (XeCl),

APPROVED BY GOVERNING BODIES:

In 2001, XTRAC[®] (PhotoMedex), a XeCl excimer laser, was cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process for the treatment of skin conditions such as vitiligo. The 510(k) clearance has subsequently been obtained for a number of targeted UVB lamps and lasers, including newer versions of the XTRAC[®] system including the XTRAC[®] Ultra, the VTRAC[™] lamp (PhotoMedex), the BClear[™] lamp (Lumenis), the 308 excimer lamp phototherapy system (Quantel Medical), MultiClear Multiwavelength Targeted Phototherapy System, Psoria-Light[™], and the Excilite[™] and Excilite μ [™] XeCl lamps. The intended use of all of these devices includes vitiligo among other dermatologic indications. Some light-emitting devices are handheld.

The oral psoralen product, methoxsalen soft gelatin capsules (previously available under the brand name Oxsoralen Ultra), has been approved by the FDA.

BENEFIT APPLICATION:

Coverage is subject to the member's specific benefits. Group-specific policy will supersede this policy when applicable.

CURRENT CODING:**CPT Codes:**

96900	Actinotherapy (ultraviolet light)
96910	Photochemotherapy; tar and ultraviolet B (Goeckerman treatment) or petrolatum and ultraviolet B
96912	; psoralens and ultraviolet A
96913	Photochemotherapy (Goeckerman and/or PUVA) for severe photoresponsive dermatoses requiring at least four to eight hours of care under direct supervision of the physician (includes application of medication and dressings)
96920	Laser treatment for inflammatory skin disease (psoriasis); total area less than 250 sq cm
96921	; 250 sq cm to 500 sq cm

96922	; over 500 sq cm
96999	Unlisted special dermatological service or procedure

HCPCS:

E0691	Ultraviolet light therapy system, includes bulbs/lamps, timer and eye protection; treatment area 2 square feet or less
E0692	; 4 foot panel
E0693	; 6 foot panel
E0694	Ultraviolet multidirectional light therapy system in six foot cabinet, includes bulbs/lamps, timer and eye protection

REFERENCES:

1. Almutawa F, Alnomair N, Wang Y et al. Systematic review of UV-based therapy for psoriasis. *Am J Clin Dermatol* 2013; 14(2):87-109.
2. Almutawa F, Thalib L, Hekman D et al. Efficacy of localized phototherapy and photodynamic therapy for psoriasis: a systematic review and meta-analysis. *Photodermatol Photoimmunol Photomed*. Jan 2015; 31(1):5-14.
3. American Academy of Dermatology. PSORIASIS CLINICAL GUIDELINE. 2021; www.aad.org/member/clinical-quality/guidelines/psoriasis
4. Amirnia M, Khodaeiani E, Fouladi RF, et al. Topical steroids versus PUVA therapy in moderate plaque psoriasis: a clinical trial along with cost analysis. *J Dermatolog Treat*. Apr 2012; 23(2):109-111.
5. Amornpinyokelt N, Asawanonda P. 8-Methoxypsoralen cream plus targeted narrowband ultraviolet B for psoriasis. *Photodermatol Photoimmunol Photomed*. Dec 2006; 22(6):285-289.
6. Archier E, Devaux S, Castela E et al. Efficacy of psoralen UV-A therapy vs. narrowband UV-B therapy in chronic plaque psoriasis: a systematic literature review. *J Eur Acad Dermatol Venereol* 2012; 26 Suppl 3:11-21.
7. Asawanonda P, et al. 308-nm excimer laser for the treatment of psoriasis; a dose-response study. *Arch Dermatol* 2000; 126:619-624.
8. Bae JM, Jung HM, Hong BY, et al. Phototherapy for vitiligo: A Systematic Review and Meta-analysis. *JAMA Dermatol*. 2017 Jul 01; 153(7):666-674.
9. Bansal S, Sahoo B, Garg V. Psoralen-narrowband UVB phototherapy in treatment of vitiligo in comparison to narrowband UVB phototherapy. *Photodermatol Photoimmunol Photomed* 2013.
10. Bonis B, et al. 308-nm UVB excimer laser for psoriasis. *Lancet* 1999; 350, 9090.
11. Callen JP, Krueger GG, Lebwohl M, McBurney EI, et al. AAD consensus statement on psoriasis therapies. *J American Acad Dermatology*, November 2003; 49(5): 897-899.
12. Cecil Textbook of Medicine, 22nd edition. Goldman Lee and Ausiella Dennis, editors. Saunders Publishing.
13. Chauhan PS, Kaur I, Dogra S, et al. Narrowband ultraviolet B versus psoralen plus ultraviolet-A therapy for severe plaque psoriasis: an Indian perspective. *Clin Exp Dermatol*. Mar 2011; 36(2):169-173.
14. Chen X, Yang M, Cheng Y et al. Narrow-band ultraviolet B phototherapy versus broadband ultraviolet B or psoralen-ultraviolet A photochemotherapy for psoriasis. *Cochrane Database Syst Rev* 2013; 10:CD009481.
15. Dayal S, Mayanka, Jain VK. Comparative evaluation of NB-UVB phototherapy and PUVA photochemotherapy in chronic plaque psoriasis. *Indian J Dermatol Venereol Leprol*. 2010; 76(5):533-537.
16. Dong DK, Pan ZY, Zhang J, et al. Efficacy and Safety of Targeted High-Intensity Medium-Band (304-312 nm) ultraviolet B light in pediatric vitiligo. *Pediatr Dermatol*. 2017 May; 34(3):266-270.
17. Elmetts CA, Lim HW, Stoff B et al. Joint American Academy of Dermatology-National Psoriasis Foundation guidelines of care for the management and treatment of psoriasis with phototherapy. *J. Am. Acad. Dermatol*. 2019 Sep;81(3):775-804.

18. El-Mofty M, Mostafa WZ, Yousef R, et al. Broadband ultraviolet-A in the treatment of psoriasis vulgaris: a randomized controlled trial. *Int J Dermatol*. Sep 2014; 53(9):1157-1164.
19. El-Zawahry BM, Bassiouny DA, Sobhi RM et al. A comparative study on efficacy of UVA1 vs. narrow-band UVB phototherapy in the treatment of vitiligo. *Photodermatol Photoimmunol Photomed* 2012; 28(2):84-90.
20. Fa Y, Lin Y, Chi XJ, et al. Treatment of vitiligo with 308-nm excimer laser: our experience from a 2-year follow-up of 979 Chinese patients. *J Eur Acad Dermatol Venereol*. Feb 2017; 31(2):337-340.
21. Feldman Steven R, Mellen Beverly G, Housman Tamara Salam, et al. Efficacy of the 308-nm excimer laser for treatment of psoriasis: results of a multicenter study. *Journal of the American Academy of Dermatology*, 2002 Jun;46(6):900-906.
22. Finlay AY. Current severe psoriasis and the rule of tens. *Br J Dermatol* 2005 May; 152(5):861-867.
23. Gerber W, Arheilger B, Ha TA, et al. Ultraviolet B 308-nm excimer laser treatment of psoriasis: a new phototherapeutic approach. *Br J Dermatol*, 2003 Dec; 149(6): 1250-1258.
24. Goldinger SM, Dummer R, Schmid P, et al. Excimer laser versus narrow-band UVB (311 nm) in the treatment of psoriasis vulgaris. *Dermatology* 2006; 213(2): 134-139.
25. Hamzavi I and Lui H. Using light in dermatology: An update on lasers, ultraviolet phototherapy, and photodynamic therapy. *Dermatol Clin*, 2005 Apr; 23(2): 199-207.
26. Ibbotson SH, Bilslund D, Cox NH, Dawe RS, et al. An update and guidance on narrowband ultraviolet B phototherapy: A British Photodermatology Group Workshop Report. *Br J Dermatology*, 2004 Au; 151(2): 283-297.
27. IOM (Institute of Medicine). 2011. *Clinical Practice Guidelines We Can Trust*. Washington, DC: The National Academies Press.
28. Khosravi H, Siegel MP, Van Voorhees AS, et al. Treatment of inverse/intertriginous psoriasis: Updated Guidelines from the Medical Board of the National Psoriasis Foundation. *J Drugs Dermatol*. Aug 01 2017;16(8):760-766.
29. Kollner K, Wimmershoff MB, Hintz C, et al. Comparison of the 308-nm excimer laser and a 308-nm excimer lamp with 311-nm narrowband ultraviolet B in the treatment of psoriasis. *Br J Dermatology*, 2005 Apr; 152(4): 750-754.
30. Kreuter A, Breuckmann F, Uhle A, et al. Low-dose UVA1 phototherapy in systemic sclerosis: Effects on acrosclerosis. *J Am Acad Dermatol*, Vol. 50, No. 5, pp. 740-747.
31. Lee E, Koo J and Berger T. UVB phototherapy and skin cancer risk: A review of the literature. *Int J Dermatol*, 2005 May; 44(5): 355-360.
32. Levin AA, Aleissa S, Dumont N, et al. A randomized, prospective, sham-controlled study of localized narrow-band UVB phototherapy in the treatment of plaque psoriasis. *J Drugs Dermatol*. 2014 Aug; 13(8):922-926.
33. Li Y, Cao Z, Guo J, Li Q, Zhu W, Kuang Y, Chen X. Assessment of efficacy and safety of UV-based therapy for psoriasis: a network meta-analysis of randomized controlled trials. *Ann Med*. 2022 Dec;54(1):159-169.
34. Lopes C, Trevisani VF, Melnik T. Efficacy and Safety of 308-nm Monochromatic Excimer Lamp versus Other Phototherapy Devices for Vitiligo: A Systematic Review with Meta-Analysis. *Am J Clin Dermatol*. 2016 Feb; 17(1):23-32.

35. Menter A, Cordoro KM, Davis DMR, et al. Joint American Academy of Dermatology-National Psoriasis Foundation guidelines of care for the management and treatment of psoriasis in pediatric patients. *J Am Acad Dermatol.* 2020 Jan; 82(1): 161-201.
36. Menter A, Korman NJ, Elmets CA et al. Guidelines of care for the management of psoriasis and psoriatic arthritis: Section 5. Guidelines of care for the treatment of psoriasis with phototherapy and photochemotherapy. *J Am Acad Dermatol* 2010; 62(1):114-135.
37. Mudigonda T, Dabade TS, Feldman SR. A review of targeted ultraviolet B phototherapy for psoriasis. *J Am Acad Dermatol.* 2012 Apr; 66(4):664-672.
38. Mudigonda T, Dabade TS, West CE et al. Therapeutic modalities for localized psoriasis: 308-nm UVB excimer laser versus nontargeted phototherapy. *Cutis* 2012 Sep; 90(3):149-154.
39. Neumann NJ, Mahnke N, Korpusik D, et al. Treatment of palmoplantar psoriasis with monochromatic excimer light (308-nm) versus cream PUVA. *Acta Derm Venereol* 2006; 86(1):22-24.
40. Nistico S, Chiricozzi A, Saraceno R et al. Vitiligo treatment with monochromatic excimer light and tacrolimus: results of an open randomized controlled study. *Photomed Laser Surg* 2012; 30(1):26-30.
41. Nistico SP, Saraceno R, Stefanescu S and Chimenti S. A 308-nm monochromatic excimer light in the treatment of palmoplantar psoriasis. *J Eur Acad Dermatol Venereol*, May 2006; 20(5): 523-526.
42. Njoo MD, Spuls PI, Bos JD, et al. Nonsurgical repigmentation therapies in vitiligo. Meta-analysis of the literature. *Arch Dermatol.* 1998 Dec; 134(12): 1532-1540.
43. Nolan BV, Yentzer BA, Feldman SR. A review of home phototherapy for psoriasis. *Dermatol Online J.* 2010 Feb; 16(2):1.
44. Oh SH, Kim T, Jee H, et al. Combination treatment of non-segmental vitiligo with a 308-nm xenon chloride excimer laser and topical high-concentration tacalcitol: a prospective, single-blinded, paired, comparative study. *J Am Acad Dermatol.* 2011 Aug; 65(2): 428-430.
45. PhotoMedex. How the XTRAC Laser Excimer Works. 2024. www.photomedex.com/pages/howwork.html
46. Poolsuwan P, Churee C, Pattamadilok B. Comparative efficacy between localized 308-nm excimer light and targeted 311-nm narrowband ultraviolet B phototherapy in vitiligo: A randomized, single-blind comparison study. *Photodermatol Photoimmunol Photomed.* Oct 12 2020.
47. Poolsuwan P, Churee C, Pattamadilok B. Comparative efficacy between localized 308-nm excimer light and targeted 311-nm narrowband ultraviolet B phototherapy in vitiligo: A randomized, single-blind comparison study. *Photodermatol Photoimmunol Photomed.* Mar 2021; 37(2): 123-130.
48. Prussick R, Wu JJ, Armstrong AW, et al. Psoriasis in solid organ transplant patients: best practice recommendations from The Medical Board of the National Psoriasis Foundation. *J Dermatolog Treat.* 2018 Jun;29(4):329-333.
49. Rodewald EJ, Housman TS, Mellen BG and Feldman SR. The efficacy of 308nm laser treatment of psoriasis compared to historical controls. *Dermatol Online J.* 2001 Dec;7(2):4.

50. Rodrigues M, Ezzedine K, Hamzavi I, et al. Current and emerging treatments for vitiligo. *J Am Acad Dermatol*. Jul 2017; 77(1):17-29.
51. Saraceno R, Nisticò SP, Capriotti E, et al. Monochromatic excimer light 308 nm in monotherapy and combined with topical khellin 4% in the treatment of vitiligo: a controlled study. *Dermatol Ther*. 2009; Jul-Aug;22(4): 391-394.
52. Seneschal J, Speeckaert R, Taïeb A, et al. Worldwide expert recommendations for the diagnosis and management of vitiligo: Position statement from the international Vitiligo Task Force-Part 2: Specific treatment recommendations. *J Eur Acad Dermatol Venereol*. 2023 Sep 15.
53. Sezer E, Erbil AH, Kurumlu Z et al. Comparison of the efficacy of local narrowband ultraviolet B (NB-UVB) phototherapy versus psoralen plus ultraviolet A (PUVA) paint for palmoplantar psoriasis. *J Dermatol* 2007 Jul; 34(7):435-440.
54. Shi Q, Li K, Fu J et al. Comparison of the 308-nm excimer laser with the 308-nm excimer lamp in the treatment of vitiligo--a randomized bilateral comparison study. *Photodermatol Photoimmunol Photomed* 2013; 29(1):27-33.
55. Sivanesan SP, Gattu S, Hong J, et al. Randomized, double-blind, placebo-controlled evaluation of the efficacy of oral psoralen plus ultraviolet A for the treatment of plaque-type psoriasis using the Psoriasis Area Severity Index score (improvement of 75% or greater) at 12 weeks. *J Am Acad Dermatol*. 2009 Nov; 61(5):793-798.
56. Spencer JM, Nossa R and Ajmeri J. Treatment of vitiligo with the 308-nm excimer laser: A pilot study. *J Am Academy of Dermatology*, 2002 May; 46(5): 727-731.
57. Sun Y, Wu Y, Xiao B, et al. Treatment of 308-nm excimer laser on vitiligo: A systemic review of randomized controlled trials. *J Dermatolog Treat*. 2015; 26(4): 347-353.
58. Taieb A, Alomar A, Bohm M et al. Guidelines for the management of vitiligo: the European Dermatology Forum consensus. *Br J Dermatol* 2013; 168(1):5-19.
59. Taneja A, Treham M and Taylor CR. 308-nm excimer laser for the treatment of psoriasis: Induration-based dosimetry. *Arch Dermatol*, 2003 Jun; 139(6): 759-764.
60. Taylor CR and Racette AL. A 308-nm excimer laser for the treatment of scalp psoriasis. *Lasers Surg Med* 2004; 34(2): 136-140.
61. Thind A, Vinay K, Mehta H, et al. Whole-body and targeted narrowband ultraviolet B phototherapy effectively stabilize acral vitiligo with negligible repigmentation beyond wrists and ankles: Results from a split-body randomized controlled trial. *Photodermatol Photoimmunol Photomed*. Mar 2024; 40(2): e12960.
62. Trehan Manju and Taylor Charles. High-dose 308-nm excimer laser for the treatment of psoriasis. *Journal for the American Academy of Dermatology*, 2002 May; 46(5):732-737.
63. Whitton ME, Pinart M, Batchelor J, et al. Interventions for vitiligo. *Cochrane Database Syst Rev*. Feb 24 2015; (2): CD003263.
64. Wollina U, Koch A, Scheibe A et al. Targeted 307 nm UVB-phototherapy in psoriasis. A pilot study comparing a 307 nm excimer light with topical dithranol. *Skin Res Technol* 2012 May; 18(2):212-218.
65. Wu Y, Sun Y, Qiu L, et al. A multicentre, randomized, split face and/or neck comparison of 308-nm excimer laser and 0.1%tacrolimus ointment for stable vitiligo plus intramuscular slow-releasing betamethasone for active vitiligo. *Br J Dermatol*. Jul2019; 181(1): 210-211.
66. www.uptodate.com/online/content/topic.do?topicKey=sclerode/2411&view=print.

67. Yang YS, Cho HR, Ryou JH, et al. Clinical study of repigmentation patterns with either narrow-band ultraviolet B (NBUVB) or 308 nm excimer laser treatment in Korean vitiligo patients. *Int J Dermatol*. Mar 2010; 49(3): 317-23.
68. Yones SS, Palmer RA, Garibaldinos TM, et al. Randomized double-blind trial of treatment of vitiligo: efficacy of psoralenUV-A therapy vs Narrowband-UV-B therapy. *Arch Dermatol*. May 2007; 143(5): 578-84.
69. Zhang C, Zhou L, Huang J, et al. A combination of Yiqiqubai granule and 308-nm excimer laser in treatment of segmental vitiligo: a prospective study of 233 patients. *J Dermatolog Treat*. Mar 21 2017:1-4.

POLICY HISTORY:

Adopted for Blue Advantage, March 2007

Available for comment April 30-June 13, 2007

Medical Policy Group, June 2007

Available for comment July 5-August 18, 2007

Medical Policy Group, May 2009

Available for comment May 16-June 28, 2009

Medical Policy Group, July 2009

Available for comment July 20-September 2, 2009

Medical Policy Group, November 2011

Medical Policy Group, December 2011

Medical Policy Group, April 2013

Medical Policy Group, April 2014

Medical Policy Group, June 2015

Medical Policy Group, January 2016

Medical Policy Group, April 2016

Available for comment April 5 through May 20, 2016

Medical Policy Group, December 2016

Medical Policy Group, August 2017

Medical Policy Group, December 2017

Medical Policy Group, April 2019

Medical Policy Group, December 2019

Medical Policy Group, March 2020: Updates to Coding Section to include CPT codes 96920, 96921, 96922, and 96999.

Medical Policy Group, January 2021

Medical Policy Group, January 2022

Medical Policy Group, December 2022

UM Committee, December 2023: Policy approved by UM Committee for use for Blue Advantage business.

Medical Policy Group, January 2024

UM Committee, January 2024: Annual review of policy approved by UM Committee for use for Blue Advantage business.

Medical Policy Group, January 2025

UM Committee January 2025: Annual review of policy approved by UM Committee for use for Blue Advantage business.

This medical policy is not an authorization, certification, explanation of benefits, or a contract. Eligibility and benefits are determined on a case-by-case basis according to the terms of the member's plan in effect as of the date services are rendered. All medical policies are based on (i) research of current medical literature and (ii) review of common medical practices in the treatment and diagnosis of disease as of the date hereof. Physicians and other providers are solely responsible for all aspects of medical care and treatment, including the type, quality, and levels of care and treatment.

This policy is intended to be used for adjudication of claims (including pre-admission certification, pre-determinations, and pre-procedure review) in Blue Cross and Blue Shield's administration of plan contracts.