



BlueCross BlueShield
of Alabama

Name of Blue Advantage Policy:
Total Artificial Hearts and Related Devices

Policy #: 033

Latest Review Date: August 2021

Category: Surgery

BACKGROUND:

Blue Advantage medical policy does not conflict with Local Coverage Determinations (LCDs), Local Medical Review Policies (LMRPs) or National Coverage Determinations (NCDs) or with coverage provisions in Medicare manuals, instructions or operational policy letters. In order to be covered by Blue Advantage the service shall be reasonable and necessary under Title XVIII of the Social Security Act, Section 1862(a)(1)(A). The service is considered reasonable and necessary if it is determined that the service is:

1. *Safe and effective;*
2. *Not experimental or investigational*;*
3. *Appropriate, including duration and frequency that is considered appropriate for the service, in terms of whether it is:*
 - *Furnished in accordance with accepted standards of medical practice for the diagnosis or treatment of the patient's condition or to improve the function of a malformed body member;*
 - *Furnished in a setting appropriate to the patient's medical needs and condition;*
 - *Ordered and furnished by qualified personnel;*
 - *One that meets, but does not exceed, the patient's medical need; and*
 - *At least as beneficial as an existing and available medically appropriate alternative.*

Routine costs of qualifying clinical trial services with dates of service on or after September 19, 2000 which meet the requirements of the Clinical Trials NCD are considered reasonable and necessary by Medicare. Providers should bill **Original Medicare for covered services that are related to **clinical trials** that meet Medicare requirements (Refer to Medicare National Coverage Determinations Manual, Chapter 1, Section 310 and Medicare Claims Processing Manual Chapter 32, Sections 69.0-69.11).*

POLICY:

Effective for dates of service on or after December 1, 2020:

For percutaneous ventricular assist devices, refer to Articles A53986 and A53988.

For ventricular assist devices, refer to NCD 20.9.1.

Blue Advantage will treat **total artificial hearts with FDA-approved devices** as a **covered benefit** when performed in a Medicare-approved heart transplant facility as a bridge to heart transplantation when ALL of the following criteria are met:

- Biventricular failure AND,
- No other reasonable medical or surgical treatment options; AND
- Are ineligible for other univentricular or biventricular support devices; AND
- Are currently listed as heart transplantation candidates

OR

- Are undergoing evaluation to determine candidacy for heart transplantation; AND
- Are not expected to survive until a donor heart can be obtained.

Blue Advantage will treat **total artificial hearts** as a **non-covered benefit** and **investigational** for all other indications, including, but not limited to, the use of total artificial hearts as destination therapy.

Blue Advantage will treat **implantable aortic counterpulsation ventricular assist devices** (e.g., the NuPulseCV iVAS and the Symphony Heart Assist System) as a **non-covered benefit** and **investigational** for all indications.

For dates of service February 26, 2018 through November 30, 2020:

For artificial hearts and related devices refer to NCD 20.9.

For percutaneous ventricular assist devices, refer to Articles A53986 & A53988.

For ventricular assist devices refer to NCD 20.9.1.

For dates of service prior to February 26, 2018:

For artificial hearts and related devices refer to NCD 20.9.

For ventricular assist devices refer to NCD 20.9.1

Blue Advantage will treat **percutaneous ventricular assist devices (pVAD)** with FDA approval or clearance as a **covered benefit** for use in patients **undergoing high risk percutaneous coronary intervention (PCI)** when All of the following are met:

- Patient has LVEF of less than 35% **AND;**

- Will undergo PCI on an unprotected left main coronary artery or last patent coronary conduit.

Blue Advantage will treat **percutaneous ventricular assist devices (pVAD)** as a **non-covered benefit** and **investigational** for all other indications.

Blue Advantage does not approve or deny procedures, services, testing, or equipment for our members. Our decisions concern coverage only. The decision of whether or not to have a certain test, treatment or procedure is one made between the physician and his/her patient. Blue Advantage administers benefits based on the members' contract and medical policies. Physicians should always exercise their best medical judgment in providing the care they feel is most appropriate for their patients. Needed care should not be delayed or refused because of a coverage determination.

DESCRIPTION OF PROCEDURE OR SERVICE:

Mechanical devices to assist or replace a failing heart have been developed over many decades of research. A ventricular assist device (VAD) is a mechanical support, attached to the native heart and vessels to augment cardiac output. The total artificial heart (TAH) replaces the native ventricles and is attached to the pulmonary artery and aorta; the native heart is typically removed. Both the VAD and TAH may be used as a bridge to heart transplantation or as destination therapy in those who are not candidates for transplantation. The VAD has also been used as a bridge to recovery in patients with reversible conditions affecting cardiac output.

Heart Failure

Heart failure may be the consequence of a number of differing etiologies, including ischemic heart disease, cardiomyopathy, congenital heart defects, or rejection of a heart transplant. The reduction of cardiac output is considered to be severe when systemic circulation cannot meet the body's needs under minimal exertion. Heart transplantation improves quality of life and has survival rates at 1-, 3-, and 5-years of 91%, 85%, and 78%, respectively. The number of candidates for transplants exceeds the supply of donor organs; thus the interest in the development of mechanical devices.

Treatment

Total Artificial Heart (TAH)

Initial research into mechanical assistance for the heart focused on the total artificial heart, a biventricular device which completely replaces the function of the diseased heart. An internal battery required frequent recharging from an external power source. Many systems utilize a percutaneous power line, but a transcutaneous power-transfer coil allows for a system without lines traversing the skin, possibly reducing the risk of infection. Because the native heart must be removed, failure of the device is synonymous with cardiac death.

A fully bioprosthetic TAH, which is fully implanted in the pericardial sac and is electrohydrolically actuated, has been developed and tested in 2 patients, but is currently experimental.

Percutaneous Ventricular Assist Devices (pVADs)

Devices in which the majority of the system's components are external to the body are for short-term use (six hours to 14 days) only, due to the increased risk of infection and need for careful, in-hospital monitoring. Some circulatory assist devices are placed percutaneously, (i.e., are not implanted). These may be referred to as percutaneous VADs (pVADs). The pVADs are placed through the femoral artery. Two different pVADs have been developed, the TandemHeart™, and the Impella® device. In the TandemHeart™ system, a catheter is introduced through the femoral vein and passed into the left atrium via transseptal puncture. Oxygenated blood is then pumped from the left atrium into the arterial system via the femoral artery. The Impella® device is introduced through a femoral artery catheter. In this device, a small pump is contained within the catheter that is placed into the left ventricle. Blood is pumped from the left ventricle, through the device, and into the ascending aorta. Adverse events associated with pVAD include access site complications such as bleeding, aneurysms, or leg ischemia. Cardiovascular complications can also occur, such as perforation, myocardial infarction (MI), stroke, and arrhythmias.

KEY POINTS:

The most recent literature search was performed for the period through April 12, 2021.

Summary of Evidence

Total Artificial Heart

For individuals who have end stage heart failure who receive total artificial hearts (TAHs) as bridge to transplant, the evidence includes case series. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, treatment-related mortality, and treatment-related morbidity. Compared with VADs, the evidence for total artificial heart in these settings is less robust. However, given the limited evidence from case series and the lack of medical or surgical options for these patients, TAH is likely to improve outcomes for a carefully selected population with end-stage biventricular heart failure awaiting transplant who are not appropriate candidates for an LVAD. The evidence is sufficient to determine qualitatively that the technology results in a meaningful improvement in the net health outcome.

For individuals who have end stage heart failure who receive TAHs as destination therapy, the evidence includes 2 case series. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, treatment-related mortality, and treatment-related morbidity. The body of evidence for TAHs as destination therapy is very limited. The evidence is insufficient to determine the effects of the technology on health outcomes.

Percutaneous Ventricular Assist Device

For individuals with cardiogenic shock or who undergo high-risk cardiac procedures who receive a pVAD, the evidence includes RCTs, observational studies, and systematic reviews. The relevant outcomes are OS, symptoms, morbid events, functional outcomes, QOL, and treatment-related mortality and morbidity. Four RCTs of pVAD vs IABP for patients in cardiogenic shock failed to demonstrate a mortality benefit and reported higher complication rates with pVAD use. Comparative observational studies were consistent with the RCT evidence. RCTs, controlled and uncontrolled observational studies, and systematic reviews of

these studies have not demonstrated a benefit of pVAD used as ancillary support for patients undergoing high-risk cardiac procedures. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals with cardiogenic shock refractory to IABP therapy who receive a pVAD, the evidence includes case series. The relevant outcomes are OS, symptoms, morbid events, functional outcomes, QOL, and treatment-related mortality and morbidity. Case series of patients with cardiogenic shock refractory to IABP have reported improved hemodynamic parameters following pVAD placement. However, these uncontrolled series do not provide evidence that pVADs improve mortality, and high rates of complications have been reported with pVAD use. The evidence is insufficient to determine the effects of the technology on health outcomes.

Practice Guidelines and Position Statements

American Association for Thoracic Surgery/International Society for Heart and Lung Transplantation

In 2020, the American Association for Thoracic Surgery and the International Society for Heart and Lung Transplantation published guidelines on selected topics in mechanical circulatory support, including recommendations on the use of pVADs. The guideline authors noted, "Compared with IABP, contemporary percutaneous circulatory support devices provide a significant increase in cardiac index and mean arterial pressure; however, reported 30-day outcomes are similar."

Table 1. 2020 Guidelines on Mechanical Circulatory Support

Recommendation	COE	LOE
"Percutaneous LV to aorta pumps of appropriate size should be considered for cardiogenic shock from primary LV failure."	IIA	B

COE: class of evidence; LOE: level of evidence; LV: left ventricular.

Society for Cardiovascular Angiography and Interventions et al

In 2015, the Society for Cardiovascular Angiography and Interventions (SCAI), the Heart Failure Society of America (HFSA), the Society of Thoracic Surgeons (STS), the American Heart Association (AHA), and the American College of Cardiology (ACC) published a clinical expert consensus statement on the use of percutaneous mechanical circulatory support (MCS) devices in cardiovascular care. This statement addressed intra-aortic balloon pumps (IABPs), left atrial (LA)-to-aorta assist device (e.g., TandemHeart®), left ventricle (LV)-to-aorta assist devices (e.g., Impella®), extracorporeal membrane oxygenation (ECMO), and methods of right-sided support. Specific recommendations were not made, but the statement reviews the use of MCS in patients undergoing high-risk percutaneous intervention, those with cardiogenic shock, and those with acute decompensated heart failure:

1. “Percutaneous MCS provides superior hemodynamic support compared to pharmacologic therapy. This is particularly apparent for the Impella[®] and TandemHeart[®] devices. These devices should remain available clinically and be appropriately reimbursed.
2. Patients in cardiogenic shock represent an extremely high risk group in whom mortality has remained high despite revascularization and pharmacologic therapies. Early placement of an appropriate MCS may be considered in those who fail to stabilize or show signs of improvement quickly after initial interventions.
3. MCS may be considered for patients undergoing high-risk PCI, such as those requiring multivessel, left main, or last patent conduit interventions, particularly if the patient is inoperable or has severely decreased ejection fraction or elevated cardiac filling pressures.”

American College of Cardiology Foundation et al

The American College of Cardiology Foundation, American Heart Association (AHA), and Heart Failure Society of American (2017) published a focused update of the 2013 recommendations released by the American College of Cardiology Foundation and AHA. Left ventricular assist device was one of several treatment options recommended for patients with refractory New York Heart Association class III or IV heart failure (stage D). If symptoms were not improved after guidelines directed management and therapy, which included pharmacologic therapy, surgical management and/or other devices, then left ventricular assist device would be an additional treatment option.

The 2017 update focused on changes in sections regarding biomarkers, comorbidities, and prevention of heart failure, while many of the previous recommendations remained unchanged. The American College of Cardiology Foundation and AHA (2013) released guidelines for the management of heart failure that included recommendations related to the use of MCS, including both durable and nondurable MCS devices. The guidelines categorized percutaneous ventricular assist devices (pVADs) and extracorporeal VADs as nondurable MCS devices.

American College of Cardiology and American Heart Association

The American College of Cardiology and American Heart Association (ACC/AHA) released guidelines for the management of heart failure in October 2013 that include recommendations related to the use of for mechanical circulatory support (MCS), including both durable and nondurable MCS devices. The guidelines categorize pVADs and extracorporeal VADs as nondurable MCS devices. The following class IIA guidelines are made related to MCS devices:

- MCS is beneficial in carefully selected patients with stage D heart failure with reduced ejection fraction (HFrEF) in whom definitive management (e.g., cardiac transplantation) or cardiac recovery is anticipated or planned. (Level of Evidence: B)
- Nondurable MCS, including the use of percutaneous and extracorporeal VADs, is reasonable as a “bridge to recovery” or “bridge to decision” for carefully selected patients with HFrEF with acute, profound hemodynamic compromise. (Level of Evidence; B)
- Durable MCS is reasonable to prolong survival for carefully selected patients with stage D HFrEF. (Level of Evidence: B)

The AHA/ACC guidelines note:

“Although optimal patient selection for MCS remains an active area of investigation, general indications for referral for MCS therapy include patients with LVEF <25% and NYHA Class III–IV functional status despite GDMT, including, when indicated, CRT, with either high predicted one to two year mortality (e.g., as suggested by markedly reduced peak oxygen consumption and clinical prognostic scores) or dependence on continuous parenteral inotropic support. Patient selection requires a multidisciplinary team of experienced advanced HF and transplantation cardiologists, cardiothoracic surgeons, nurses, and ideally, social workers and palliative care clinicians.”

In 2012, AHA published recommendations for the use of MCS. These guidelines define nondurable MCS as intra-aortic balloon pump (IABP), extracorporeal membrane oxygenation, extracorporeal VADs, and pVADs. The following recommendations were made regarding indications for use of MCS, including durable and nondurable devices:

- MCS for bridge-to-transplant indication should be considered for transplant-eligible patients with end-stage heart failure who are failing optimal medical, surgical, and/or device therapies and at high risk of dying before receiving a heart transplantation. (Class I; Level of Evidence B).
- Implantation of MCS in patients before the development of advanced heart failure... is associated with better outcomes. Therefore, early referral of heart failure patients is reasonable. (Class IIa; Level of Evidence B).
- MCS with a durable, implantable device for permanent therapy or destination therapy is beneficial for patients with advanced heart failure, high one year mortality resulting from heart failure, and the absence of other life-limiting organ dysfunction; who are failing medical, surgical, and/or device therapies; and who are ineligible for heart transplantation. (Class I; Level of Evidence B).
- Elective rather than urgent implantation of destination therapy can be beneficial when performed after optimization of medical therapy in advanced heart failure patients who are failing medical, surgical, and/or device therapies. (Class IIa; Level of Evidence C).
- Urgent nondurable MCS is reasonable in hemodynamically compromised heart failure patients with end-organ dysfunction and/or relative contraindications to heart transplantation/durable MCS that are expected to improve with time and restoration of an improved hemodynamic profile. (Class IIa; Level of Evidence C).
- These patients should be referred to a center with expertise in the management of durable MCS and patients with advanced heart failure. (Class I; Level of Evidence C).
- Patients who are ineligible for heart transplantation because of pulmonary hypertension related to heart failure alone should be considered for bridge to potential transplant eligibility with durable, long-term MCS. (Class IIa; Level of Evidence B).

Heart Failure Society of America

The Heart Failure Society of America published guidelines in 2010 on surgical approaches to the treatment of heart failure. The following recommendations were made regarding left ventricular assist devices:

- Patients awaiting heart transplantation who have become refractory to all means of medical circulatory support should be considered for a mechanical support device as a bridge to transplant. (Strength of Evidence = B)
- Permanent mechanical assistance using an implantable assist device may be considered in highly selected patients with severe HF [heart failure] refractory to conventional therapy who are not candidates for heart transplantation, particularly those who cannot be weaned from intravenous inotropic support at an experienced HF center. (Strength of Evidence = B)
- Patients with refractory HF and hemodynamic instability, and/or compromised end-organ function, with relative contraindications to cardiac transplantation or permanent mechanical circulatory assistance expected to improve with time or restoration of an improved hemodynamic profile should be considered for urgent mechanical circulatory support as a "bridge to decision." These patients should be referred to a center with expertise in the management of patients with advanced HF. (Strength of Evidence = C)

U.S. Preventive Services Task Force Recommendations

Not applicable

KEY WORDS:

Ventricular assist device, biventricular support, BIVAD, cardiac support, heart transplantation (transplant), LVAD, VAD, destination therapy, HeartWare®, Impella LV®, Impella 2.5, Impella 2.5 circulatory assist device, DeBakey, percutaneous ventricular assist device, pVAD, TandemHeart®, Berlin Heart EXCOR®, Impella RP, Carmat, bioprosthetic artificial heart, HeartMate III™, Total Artificial Heart, TAH, CardioWest™ Total Artificial Heart, HeartMate II®, SynCardia artificial heart, Right Ventricular Assist Device, RVAD, PediMag®, short-term continuous flow ventricular assist devices, STCF-VADs, intraluminal axial support, HeartAssist 5 Pediatric Ventricular Assist Device; NuPulseCV iVAS; Symphony Heart Assist System; CentriMag® Blood Pump; implantable aortic counterpulsation ventricular assist devices; Intravascular Ventricular Assist Systems; iVAS); C-Pulse, CardioVAD

APPROVED BY GOVERNING BODIES:

A number of mechanical circulatory support devices have received approval or clearance for marketing by FDA. These devices are summarized in Table 2, and described further in the sections below.

Table 2: Available Mechanical Circulatory Support Devices

Device	Manufacturer	Approval Date	FDA Clearance	PMA, HDE, or 510(k) No.	Indication
--------	--------------	---------------	---------------	-------------------------	------------

Thoratec® IVAD	Thoratec	Aug 2004	PMA supplement	P870072	Bridge to transplant and postcardiotomy
DeBakey VAD® Child	MicroMed	Feb 2004	HDE	H030003	Bridge to transplant in children 5-16 years of age
HeartMate II®	Thoratec	Apr 2008	PMA	P060040	Bridge to transplant and destination therapy
Centrimag®	Levitronix	Oct 2008	HDE	H070004	Postcardiotomy
Berlin Heart EXCOR® Pediatric VAD	Berlin	Dec 2011	HDE	H100004	Bridge to transplant
HeartWare® Ventricular Assist System	HeartWare	Dec 2012	PMA	P100047	Bridge to transplant, and destination therapy
HeartMate III™ Left Ventricular Assist System	Thoratec	Aug 2017	PMA	P160054 P160054/S008	Bridge to transplant, and destination therapy

FDA: Food and Drug Administration; HDE: humanitarian device exemption; PMA: premarket approval

Ventricular Assist Devices

In December 1995, the Thoratec® Ventricular Assist Device System (Thoratec Corp., Pleasanton, CA) was approved by the FDA through the premarket approval process for use as a bridge to transplantation in patients suffering from end-stage heart failure. The patient should meet all of the following criteria:

- candidate for cardiac transplantation,
- imminent risk of dying before donor heart procurement, and
- dependence on, or incomplete response to, continuous vasopressor support.

In May 1998, supplemental approval for the above device was given for the indication for postcardiotomy patients who are unable to be weaned from cardiopulmonary bypass. In June 2001, supplemental approval was given for a portable external driver to permit excursions within a 2-hour travel radius of the hospital in the company of a trained caregiver. In November 2003, supplemental approval was given to market the device as Thoratec® Paracorporeal VAD. In August 2004, supplemental approval was given to a modified device to be marketed as the Thoratec® Implantable VAD for the same indications. In January 2008, supplemental approval was given to delete Paracorporeal VAD use.

In February 2004, the FDA approved the DeBakey VAD® Child under the HDE approval process. According to the FDA, this device is indicated under HDE for both home and hospital use for children who are between ages 5 and 16 years and who have end-stage ventricular failure requiring temporary mechanical blood circulation until a heart transplant is performed. In April 2008, continuous flow device HeartMate II® LVAS (Thoratec, Pleasanton, CA) was approved by the FDA through the premarket approval process for use as a bridge to transplantation in cardiac transplant candidates at risk of imminent death from nonreversible left ventricular failure. The Heartmate II® LVAS is intended for use both inside and outside the hospital. In January 2010, the device received the added indication as destination therapy for use in patients with New York Heart Association (NYHA) Class IIIB or IV end-stage left ventricular failure who have received optimal medical therapy for at least 45 of the last 60 days and are not candidates for cardiac transplantation.

In October 2008, device Centrimag® Right Ventricular Assist Device (Levitronix, Zurich) was approved by the FDA under the HDE to provide temporary circulatory support for up to 14 days for patients in cardiogenic shock due to acute right-sided heart failure.

In December 2011, the Berlin Heart EXCOR® Pediatric VAD was approved via HDE. The indications for this device are pediatric patients with severe isolated left ventricular or biventricular dysfunction who are candidates for cardiac transplant and require circulatory support.

In December 2012, device HeartWare® Ventricular Assist System (HeartWare, Inc., Framingham, Mass.) was approved by the FDA using the INTERMACS registry as a control. INTERMACS registry was established in 2005 as a joint effort involving the FDA, National Heart, Lung and Blood Institute (NHLBI), Centers for Medicare and Medicaid Services (CMS), clinicians, scientists, and industry. This was the first time the FDA approved an LVAD using registry data as a control. INTERMACS is managed by the University of Alabama at Birmingham.

In August 2016, HeartWare® recalled its VAD Pumps due to a design flaw that was deemed by FDA as potentially causing serious injuries or death (class I recall). The devices affected were manufactured and distributed from March 2006 and May 2018. FDA product codes 204 and 017.

In September 2017, HeartWare® Ventricular Assist System (HeartWare, Inc., Framingham, Mass.) was approved by the FDA for providing long-term hemodynamic support (e.g., destination therapy) in patients with advanced heart failure.

In August 2017, the HeartMate™ 3 Left Ventricular Assist System (Thoratec Corp., Pleasanton, CA) was approved by the FDA for providing short-term hemodynamic support (e.g., bridge to transplant or bridge to myocardial recovery) in patients with advanced refractory left ventricular heart failure.

In October 2018, the HeartMate™ 3 Left Ventricular Assist System (Thoratec Corp., Pleasanton, CA) was approved by the FDA for providing long-term hemodynamic support (e.g., destination therapy) in patients with advanced heart failure.

A class I recall was issued for the HeartMate 3™ in April 2018 affecting all manufacturing dates. FDA product code: DSQ.

Total Artificial Heart

In 2004, the temporary CardioWest™ Total Artificial Heart (SynCardia Systems) was approved by FDA through the premarket approval process for use as a bridge to transplant in cardiac transplant-eligible candidates at risk of imminent death from biventricular failure. This device is also intended for use inside the hospital. In 2010, FDA approved a name change to SynCardia Temporary Total Artificial Heart. FDA product code: LOZ.

In 2006, the AbioCor® Implantable Replacement Heart System (Abiomed) was approved by FDA through the humanitarian device exemption (H040006) process in severe biventricular end-stage heart disease patients who are not cardiac transplant candidates and who:

- are younger than 75 years of age;
- require multiple inotropic support;
- are not treatable by left VAD destination therapy; and
- are not weanable from biventricular support if on such support.

In addition to meeting other criteria, patients who are candidates for the AbioCor® TAH must undergo a screening process to determine if their chest volume is large enough to hold the device. The device is too large for about 90% of women and for many men.

****NOTE: The Carmat bioprosthetic total artificial heart has not been FDA approved.**

Percutaneous VADs (Circulatory Assist Devices)

Table 3. Available Mechanical Circulatory Support Devices

Device	Manufacturer	Approval Date	FDA Clearance	PMA, 510(k) No.	Indication
TandemHeart®	Cardiac Assist	Sep 2005	510(k)	K110493	Temporary left ventricular bypass of ≤6 h

Impella® Recover LP 2.5	Abiomed	May 2008	510(k)	K063723	Partial circulatory support using extracorporeal bypass control unit for ≤6 h
Impella® 2.5 System	Abiomed	Mar 2015	PMA	P140003	Temporary ventricular support for ≤6 h

FDA: U.S. Food and Drug Administration; PMA: premarket approval.

Comparative Efficacy of Left VAD Devices

The mechanism of operation of left VADs has changed since their introduction. The earliest devices were pulsatile positive displacement pumps. These pumps have been largely replaced by axial continuous-flow pumps. More recently centrifugal continuous-flow pumps have also been introduced.

The evidence of the comparative efficacy of centrifugal continuous-flow vs axial continuous-flow devices consists of 2 randomized controlled trials of 2 different centrifugal continuous-flow devices. The MOMENTUM 3 trial compared HeartMate III™ centrifugal continuous-flow device with the HeartMate II® axial continuous-flow device in patients indicated for circulatory support as a bridge to transplant or destination therapy. HeartMate III™ received PMA approval in August 2017 but was recalled in April 2018. The ENDURANCE trial compared HeartWare® centrifugal continuous-flow device with the HeartMate II® axial continuous-flow device in patients indicated for circulatory support as destination therapy. HeartWare® is FDA-approved as a bridge to transplantation device. Both trials found the centrifugal device to be noninferior to the axial device for the primary, composite outcome including measures of survival, freedom from disabling stroke, and freedom from device failure. While there are fewer device failures with the centrifugal devices without a significant increase in disabling stroke, the HeartWare® device was associated with increased risk of any stroke over a period of 2 years.

The evidence on the comparative efficacy of continuous-flow vs pulsatile-flow devices consists of a randomized controlled trial and several nonrandomized comparative studies. The randomized controlled trial reported fairly large differences in a composite outcome measure favoring the continuous-flow devices, with increases in revision and reoperation rates for the pulsatile device group being the largest factor driving the difference in outcomes. Other nonrandomized comparative studies, including a database study with large numbers of patients, have not reported important differences in clinical outcomes between devices.

Percutaneous Ventricular Assist Devices (circulatory assist devices)

The Impella® Recover LP 2.5 Percutaneous Cardiac Support System (Abiomed, Aachen, Germany) received FDA 510(k) approval in May 2008 for short-term (less than six hours) use in patients requiring circulatory support.

In March 2015, the Impella® 2.5 System received approval through the PMA process for temporary ventricular support during high-risk percutaneous coronary interventions.

The TandemHeart® (Cardiac Assist, Pittsburgh) received a similar 510(k) approval for short-term circulatory support in September 2005.

Several other devices are in clinical trials or awaiting FDA review.

BENEFIT APPLICATION:

Coverage is subject to member’s specific benefits. Group specific policy will supersede this policy when applicable.

CURRENT CODING:

CPT codes:

33927	Implantation of a total replacement heart system (artificial heart) with recipient cardiectomy
33928	Removal and replacement of total replacement heart system (artificial heart)
33929	Removal of a total replacement heart system (artificial heart) for heart transplantation (List separately in addition to code for primary procedure)

As of 1/1/22, aortic counterpulsation ventricular assist devices should be reported using the unlisted code below.

33999	Unlisted procedure, cardiac surgery
-------	-------------------------------------

HCPCS Codes:

L8698	Miscellaneous component, supply or accessory for use with total artificial heart system
-------	---

PREVIOUS CODING:

Aortic Counterpulsation Ventricular Assist Devices

0451T	Insertion or replacement of a permanently implantable aortic counterpulsation ventricular assist system, endovascular approach, and programming of sensing and therapeutic parameters; complete system (counterpulsation device, vascular graft, implantable vascular hemostatic seal, mechano-electrical skin interface and subcutaneous electrodes) (Deleted 12/31/21)
-------	---

0452T	Insertion or replacement of a permanently implantable aortic counterpulsation ventricular assist system, endovascular approach, and programming of sensing and therapeutic parameters; aortic counterpulsation device and vascular hemostatic seal (Deleted 12/31/21)
0453T	Insertion or replacement of a permanently implantable aortic counterpulsation ventricular assist system, endovascular approach, and programming of sensing and therapeutic parameters; mechano-electrical skin interface (Deleted 12/31/21)
0454T	Insertion or replacement of a permanently implantable aortic counterpulsation ventricular assist system, endovascular approach, and programming of sensing and therapeutic parameters; subcutaneous electrode (Deleted 12/31/21)
0455T	Removal of permanently implantable aortic counterpulsation ventricular assist system; complete system (aortic counterpulsation device, vascular hemostatic seal, mechano-electrical skin interface and electrodes) (Deleted 12/31/21)
0456T	Removal of permanently implantable aortic counterpulsation ventricular assist system; aortic counterpulsation device and vascular hemostatic seal (Deleted 12/31/21)
0457T	Removal of permanently implantable aortic counterpulsation ventricular assist system; mechano-electrical skin interface (Deleted 12/31/21)
0458T	Removal of permanently implantable aortic counterpulsation ventricular assist system; subcutaneous electrode (Deleted 12/31/21)
0459T	Relocation of skin pocket with replacement of implanted aortic counterpulsation ventricular assist device, mechano-electrical skin interface and electrodes Repositioning of previously implanted aortic counterpulsation ventricular assist device; subcutaneous electrode (Deleted 12/31/21)
0460T	Repositioning of previously implanted aortic counterpulsation ventricular assist device; subcutaneous electrode (Deleted 12/31/21)
0461T	Repositioning of previously implanted aortic counterpulsation ventricular assist device; aortic counterpulsation device (Deleted 12/31/21)
0462T	Programming device evaluation (in person) with iterative adjustment of the implantable mechano-electrical skin interface and/or external driver to test the function of the device and select optimal permanent programmed values with analysis, including review and report, implantable aortic counterpulsation ventricular assist system, per day (Deleted

	12/31/21)
0463T	Interrogation device evaluation (in person) with analysis, review and report, includes connection, recording and disconnection per patient encounter, implantable aortic counterpulsation ventricular assist system, per day (Deleted 12/31/21)

REFERENCES:

1. 2008 OPTN/SRTR Annual Report 1998-2007. HHS/HRSA/HSB/DOT. Accessed 11/30/2010.
2. Aaronson KD, Eppinger MJ, Dyke DB et al. Left ventricular assist device therapy improves utilization of donor hearts. J Am Coll Cardiol 2002; 39(8):1247-54.
3. Aaronson KD, Slaughter MS, Miller LW, et al. Use of an intrapericardial, continuous-flow, centrifugal pump in patients awaiting heart transplantation. Circulation. Jun 26 2012; 125(25):3191-3200.
4. Acharya D, Loyaga-Rendon RY, Pamboukian SV, et al. Ventricular assist device in acute myocardial infarction. J Am Coll Cardiol. Apr 26 2016; 67(16):1871-1880.
5. Aissaoui N, Morshuis M, Maoulida H, et al. Management of end-stage heart failure patients with or without ventricular assist device: an observational comparison of clinical and economic outcomes. Eur J Cardiothorac Surg. 2018 53(1).
6. Alasnag MA, Gardi DO, Elder M, et al. Use of the Impella 2.5 for prophylactic circulatory support during elective high-risk percutaneous coronary intervention. Cardiovasc Revasc Med. Sep-Oct 2011; 12(5):299-303.
7. Alba AC, McDonald M, Rao V et al. The effect of ventricular assist devices on long-term post-transplant outcomes: a systematic review of observational studies. Eur J Heart Fail 2011; 13(7):785-95.
8. Almond CS, Buchholz H, Massicotte P et al. Berlin Heart EXCOR Pediatric ventricular assist device Investigational Device Exemption study: study design and rationale. Am Heart J 2011; 162(3):425-35 e6.
9. Almond CS, Morales DL, Blackstone EH, et al. Berlin Heart EXCOR pediatric ventricular assist device for bridge to heart transplantation in US children. Circulation. Apr 23 2013; 127(16):1702-1711.
10. Al-Sarie M, Rauf A, Kfoury AG, et al. Myocardial structural and functional response after long-term mechanical unloading with continuous flow left ventricular assist device: axial versus centrifugal flow. JACC Heart Fail. Jul 2016; 4(7):570-576.
11. Agrawal S, Garg L, Shah M, et al. Thirty-Day Readmissions after Left Ventricular Assist Device Implantation in the United States: Insights from the Nationwide Readmissions Database. Circ Heart Fail 2018 11(3):e004628.
12. American Medical Association. CPT 2021 Professional.
13. Arnold SV, Jones PG, Allen LA, et al. Frequency of poor outcome (death or poor quality of life) after left ventricular assist device for destination therapy: results from the INTERMACS Registry. Circ Heart Fail. Aug 2016; 9(8).

14. Aryana A, Gearoid O'Neill P, Gregory D, et al. Procedural and clinical outcomes after catheter ablation of unstable ventricular tachycardia supported by a percutaneous left ventricular assist device. *Heart Rhythm*. Jul 2014; 11(7):1122-1130.
15. Bank AJ, Mir SH, Nguyen DQ et al. Effects of left ventricular assist devices on outcomes in patients undergoing heart transplantation. *Ann Thorac Surg* 2000; 69(5):1369-74; discussion 75.
16. Basir MB, Schreiber TL, Grines CL, et al. Effect of early initiation of mechanical circulatory support on survival in cardiogenic shocks. *Am J Cardiol*. Mar 15 2017;119(6):845-851.
17. Birks EJ, George RS, Hedger M, et al. Reversal of severe heart failure with a continuous-flow left ventricular assist device and pharmacological therapy: a prospective study. *Circulation*. Feb 1 2011; 123(4):381-390.
18. Birks EJ, Tansley PD, Hardy J, et al. Left ventricular assist device and drug therapy for the reversal of heart failure. *N Engl J Med*. Nov 2 2006;355(18):1873-1884.
19. Blue Cross Blue Shield Association, Technology Evaluation Center (TEC) Assessment, January 1997, Vol. 11, No. 26.
20. Blue Cross Blue Shield Association, Technology Evaluation Center (TEC) Assessment 1996: Ventricular assist devices in bridging to heart transplantation. 1996; Volume 11;Tab 26.
21. Blue Cross Blue Shield Association. Left-ventricular assist devices as destination therapy for end-stage heart failure. Technology Evaluation Center (TEC) Assessment, December 2002, Vol. 17, No. 19.
22. Blue Cross Blue Shield Association. Ventricular assist devices and total artificial hearts. Medical Policy Reference Manual, September 2010.
23. Blume ED, Rosenthal DN, Rossano JW, et al. Outcomes of children implanted with ventricular assist devices in the United States: First analysis of the Pediatric Interagency Registry for Mechanical Circulatory Support (PediMACS). *J Heart Lung Transplant*. May 2016; 35(5):578-584.
24. Briasoulis A, Telila T, Palla M, et al. Meta-analysis of usefulness of percutaneous left ventricular assist devices for high-risk percutaneous coronary interventions. *Am J Cardiol*. Aug 1 2016;118(3):369-375.
25. Bulic A, Maeda K, Zhang Y, et al. Functional status of United States children supported with a left ventricular assist device at heart transplantation. *J Heart Lung Transplant*. Aug 2017; 36(8):890-896.
26. Burkhoff D, Cohen H, Brunckhorst C, et al. A randomized multicenter clinical study to evaluate the safety and efficacy of the TandemHeart percutaneous ventricular assist device versus conventional therapy with intra-aortic balloon pumping for treatment of cardiogenic shock. *Am Heart J*. Sep 2006; 152(3):469 e461-468.
27. Carpentier A, Latremouille C, Cholley B, et al. First clinical use of a bioprosthetic total artificial heart: report of two cases. *Lancet*. Oct 17 2015; 386(10003):1556-1563.
28. Chen S, Lin A, Liu E, et al. Outpatient outcomes of pediatric patients with left ventricular assist devices. *ASAIO J*. Mar-Apr 2016; 62(2):163-168.
29. Cheng JM, den Uil CA, Hoeks SE, et al. Percutaneous left ventricular assist devices vs. intra-aortic balloon pump counterpulsation for treatment of cardiogenic shock: a meta-analysis of controlled trials. *Eur Heart J*. Sep 2009; 30(17):2102-2108.

30. Conway J, Al-Aklabi M, Granoski D, et al. Supporting pediatric patients with short-term continuous-flow devices. *J Heart Lung Transplant*. May 2016; 35(5):603-609.
31. Copeland JG, Copeland H, Gustafson M et al. Experience with more than 100 total artificial heart implants. *J Thorac Cardiovasc Surg* 2012; 143(3):727-34.
32. Copeland JG, Smith RG, Arabia FA et al. Cardiac replacement with a total artificial heart as a bridge to transplantation. *N Engl J Med* 2004; 351(9):859-67.
33. Dangas GD, Kini AS, Sharma SK, et al. Impact of hemodynamic support with Impella 2.5 versus intra-aortic balloon pump on prognostically important clinical outcomes in patients undergoing high-risk percutaneous coronary intervention (from the PROTECT II randomized trial). *Am J Cardiol*. Jan 15 2014; 113(2):222-228.
34. Davies RR, Russo MJ, Hong KN et al. The use of mechanical circulatory support as a bridge to transplantation in pediatric patients: an analysis of the United Network for Organ Sharing database. *J Thorac Cardiovasc Surg* 2008; 135(2):421-427, 427 e421.
35. Dell'Aquila AM, Schneider SR, Stypmann J, et al. Survival results after implantation of intrapericardial third-generation centrifugal assist device: an INTERMACS-matched comparison analysis. *Artif Organs*. May 2014; 38(5):383-390.
36. De Robertis F, Birks EJ, Rogers P et al. Clinical performance with the Levitronix Centrimag short-term ventricular assist device. *J Heart Lung Transplant* 2006; 25(2):181-6.
37. De Robertis F, Rogers P, Amrani M et al. Bridge to decision using the Levitronix CentriMag short-term ventricular assist device. *J Heart Lung Transplant* 2008; 27(5):474-8.
38. Deo SV, Sung K, Daly RC, et al. Cardiac transplantation after bridged therapy with continuous flow left ventricular assist devices. *Heart Lung Circ*. Mar 2014; 23(3):224-228.
39. Dickstein K, Vardas PE, Auricchio A et al. 2010 Focused Update of ESC Guidelines on device therapy in heart failure: an update of the 2008 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure and the 2007 ESC guidelines for cardiac and resynchronization therapy. Developed with the special contribution of the Heart Failure Association and the European Heart Rhythm Association. *Eur Heart J* 2010; 31(21):2677-87.
40. Dickstein K, Cohen-Solal A, Filippatos G, et al. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2008: the Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2008 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association of the ESC (HFA) and endorsed by the European Society of Intensive Care Medicine (ESICM). *Eur Heart J*. Oct 2008; 29(19):2388-2442.
41. Dixon SR, Henriques JP, et al. A prospective feasibility trial investigating the use of the Impella 2.5 system in patients undergoing high-risk percutaneous coronary intervention (The PROTECT I Trial): Initial U.S. experience. *JACC Cardiovasc Interv*, February 2009; 2(2): 91-96.
42. Dowling RD, Gray LA, Jr., Etoch SW et al. Initial experience with the AbioCor implantable replacement heart system. *J Thorac Cardiovasc Surg* 2004; 127(1):131-41.
43. Estep JD, Starling RC, Horstmanshof DA, et al. Risk assessment and comparative effectiveness of left ventricular assist device and medical management in ambulatory

- heart failure patients: results from the ROADMAP study. *J Am Coll Cardiol*. Oct 20 2015; 66(16):1747-1761.
44. FDA. Summary of Safety and Probable Benefit - H040006: AbioCor Implantable Replacement Heart. 2006; http://www.accessdata.fda.gov/cdrh_docs/pdf4/H040006b.pdf. Accessed July 31, 2017FDA information: Debakey VAD Child. Available online at: www.accessdata.fda.gov/cdrh_docs/pdf3/H030003b.pdf. Last reviewed March 2012.
 45. Food and Drug Administration. Summary of Safety and Probable Benefit - H040006: AbioCor Implantable Replacement Heart. 2006; https://www.accessdata.fda.gov/cdrh_docs/pdf4/H040006b.pdf. Accessed August 8, 2018.
 46. FDA information: FDA approves pump for heart failure patients awaiting heart transplant. Available online at: www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm328818.htm. November 20, 2012.
 47. FDA information: Thoratec Heartmate II. Available online at: www.accessdata.fda.gov/cdrh_docs/pdf6/P060040b.pdf. Last reviewed March 2012.
 48. Feldman D, Pamboukian SV, Teuteberg JJ, et al. The 2013 International Society for Heart and Lung Transplantation Guidelines for mechanical circulatory support: executive summary. *J Heart Lung Transplant*. Feb 2013; 32(2):157-187.
 49. Fraser CD, Jr., Jaquiss RD, Rosenthal DN, et al. Prospective trial of a pediatric ventricular assist device. *N Engl J Med*. Aug 9 2012; 367(6):532-541.
 50. Frazier OH, Gemmato C, Myers TJ et al. Initial clinical experience with the HeartMate II axial-flow left ventricular assist device. *Tex Heart Inst J* 2007; 34(3):275-81.
 51. Frazier OH, Rose EA, McCarthy P et al. Improved mortality and rehabilitation of transplant candidates treated with a long-term implantable left ventricular assist system. *Ann Surg* 1995; 222(3):327-336; discussion 336-338.
 52. Frieden J. FDA panel gives okay to pediatric use of LVAD. July 2011, www.medpagetoday.com/Cardiology/HeartTransplantation/27738.
 53. Gafoor S, Franke J, Lam S, Reinartz M, Bertog S, Vaskelyte L, Hofmann I, Sievert H. Devices in heart failure--the new revolution. *Circ J*. 2015; 79(2):237-44.
 54. Geligns AC, et al. Evolving costs of long-term left ventricular assist device implantation. *Annals of Thoracic Surgery*, 1997; 64:1312-19.
 55. Griffith BP, Anderson MB, Samuels LE et al. The RECOVER I: A multicenter prospective study of Impella 5.0/LD for postcardiotomy circulatory support. *J Thorac Cardiovasc Surg*. 2013 Feb; 145(2):548-54
 56. Grimm JC, Sciortino CM, Magruder JT, et al. Outcomes in Patients Bridged With Univentricular and Biventricular Devices in the Modern Era of Heart Transplantation. *Ann Thorac Surg*. Jul 2016; 102(1):102-108.
 57. Goldstein DJ, Ox MC, and Rose EA. Implantable left ventricular assist devices. *N England J Med* 1998; 339(21):1522-1533.
 58. Heart Failure Society of America, Lindenfeld J, Albert NM, et al. HFSA 2010 Comprehensive Heart Failure Practice Guideline. *J Card Fail*. Jun 2010;16(6):e1-194.
 59. Hunt SA, Abraham WT, Chin MH et al. 2009 focused update incorporated into the ACC/AHA 2005 Guidelines for the Diagnosis and Management of Heart Failure in Adults: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines: developed in collaboration with the

- International Society for Heart and Lung Transplantation. *Circulation* 2009; 119(14):e391-479.
60. Hunt SA, Abraham WT, Chin MH et al. ACC/AHA 2005 Guideline Update for the Diagnosis and Management of Chronic Heart Failure in the Adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure): developed in collaboration with the American College of Chest Physicians and the International Society for Heart and Lung Transplantation: endorsed by the Heart Rhythm Society. *Circulation* 2005; 112(12):e154-235.
 61. Jeevanandam V, Jayakar D, Anderson AS, et al. Circulatory assistance with a permanent implantable IABP: Initial human experience. *Circulation*. 2002; 106 (12 Suppl 1):I183-I188.
 62. John R, Kamdar F, Liao K et al. Improved survival and decreasing incidence of adverse events with the HeartMate II left ventricular assist device as bridge-to-transplant therapy. *Ann Thorac Surg* 2008; 86(4):1227-34; discussion 34-5.
 63. John R., et al. Impact of current management practices on early and late date in more than 500 consecutive cardiac transplant recipients. *Annals of Surgery* 2000; 232(3):302-11.
 64. Jordan LC, Ichord RN, Reinhartz O, et al. Neurological complications and outcomes in the Berlin Heart EXCOR(R) pediatric investigational device exemption trial. *J Am Heart Assoc*. Jan 2015; 4(1):e001429.
 65. Jorde UP, Kushwaha SS, Tatoes AJ, et al. Results of the destination therapy post-food and drug administration approval study with a continuous flow left ventricular assist device: a prospective study using the INTERMACS registry (Interagency Registry for Mechanically Assisted Circulatory Support). *J Am Coll Cardiol*. May 6 2014; 63(17):1751-1757.
 66. Kaiser C. Pediatric Cardiac Assist Device Approved. *MedPage Today* 2011 www.medpagetoday.com/Cardiology/Heart_Transplantation/.
 67. Kar B, Forrester M, Gemmato C et al. Use of the TandemHeart percutaneous ventricular assist device to support patients undergoing high-risk percutaneous coronary intervention. *J Invasive Cardiol* 2006; 18(3):93-6.
 68. Kar B, Gregoric ID, Basra SS et al. The percutaneous ventricular assist device in severe refractory cardiogenic shock. *J Am Coll Cardiol* 2011; 57(6):688-96.
 69. Kato TS, Chokshi A, Singh P et al. Effects of continuous-flow versus pulsatile-flow left ventricular assist devices on myocardial unloading and remodeling. *Circ Heart Fail* 2011; 4(5):546-53.
 70. Kirklin JK, Naftel DC, Stevenson LW et al. INTERMACS database for durable devices for circulatory support: first annual report. *J Heart Lung Transplant* 2008; 27(10):1065-72.
 71. Kirklin JK, Pagani FD, Goldstein DJ, et al. American Association for Thoracic Surgery/International Society for Heart and Lung Transplantation guidelines on selected topics in mechanical circulatory support. *J Heart Lung Transplant*. Mar 2020; 39(3): 187-219.
 72. Kontogiannis CD, Malliaras K, Kapelios CJ, Mason JW, Nanas JN. Continuous internal counterpulsation as a bridge to recovery in acute and chronic heart failure. *World J Transplant*. 2016 Mar 24; 6(1):115-24.

73. Korfer R, et al. Temporary pulsatile ventricular assist devices and biventricular assist devices. *Annual of Thoracic Surgery* 1999;68(2): 678-83.
74. Koul B, et al. Temporary pulsatile ventricular assist devices as bridge to heart transplantation. *Annals of Thoracic Surgery*, January 1998, 65:1625-30.
75. Kovacic JC, Kini A, Banerjee S, et al. Patients with 3-vessel coronary artery disease and impaired ventricular function undergoing PCI with Impella 2.5 hemodynamic support have improved 90-day outcomes compared to intra-aortic balloon pump: a sub-study of the PROTECT II trial. *J Interv Cardiol*. Feb 2015; 28(1):32-40.
76. Kovacic JC, Nguyen HT, Karajgikar R, et al. The Impella Recover 2.5 and TandemHeart ventricular assist devices are safe and associated with equivalent clinical outcomes in patients undergoing high-risk percutaneous coronary intervention. *Catheter Cardiovasc Interv*. Jul 01 2013; 82(1):E28-37.
77. Lauten A, Engstrom AE, Jung C, et al. Percutaneous left-ventricular support with the Impella-2.5-assist device in acute cardiogenic shock: results of the Impella-EUROSHOCK-registry. *Circ Heart Fail*. Jan 2013; 6(1):23-30.
78. Lemaire A, Anderson MB, Lee LY, et al. The Impella device for acute mechanical circulatory support in patients in cardiogenic shock. *Ann Thorac Surg*. Jan 2014; 97(1):133-138.
79. Lemaire A, Anderson MB, Lee LY, et al. The Impella device for acute mechanical circulatory support in patients in cardiogenic shock. *Ann Thorac Surg*. Jan 2014; 97(1):133-138.
80. Lim KM, Constantino J, Gurev V et al. Comparison of the effects of continuous and pulsatile left ventricular-assist devices on ventricular unloading using a cardiac electromechanics model. *J Physiol Sci* 2012; 62(1):11-19.
81. Lindenfeld J, Albert NM, Boehmer JP et al. HFSA 2010 Comprehensive Heart Failure Practice Guideline. *J Card Fail* 2010; 16(6):e1-194.
82. Long JW, Kfoury AG, Slaughter MS et al. Long-term destination therapy with the HeartMate XVE left ventricular assist device: improved outcomes since the REMATCH study. *Congest Heart Fail* 2005; 11(3):133-8.
83. Maini B, Naidu SS, Mulukutla S, et al. Real-world use of the Impella 2.5 circulatory support system in complex high-risk percutaneous coronary intervention: the USpella Registry. *Catheter Cardiovasc Interv*. Nov 1 2012;80(5):717-725.
84. Maybaum S, Mancini D, Xydias S, et al. Cardiac improvement during mechanical circulatory support: a prospective multicenter study of the LVAD Working Group. *Circulation*. May 15 2007;115(19):2497-2505.
85. McCarthy PM, et al. One-hundred patients with the HeartMate left ventricular assist device: Evolving concepts and technology. *Journal of Thoracic and Cardiovascular Surgery*, April 1998, pp. 904-12.
86. McMurray JJ, Adamopoulos S, Anker SD, et al. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: The Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2012 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association (HFA) of the ESC. *Eur Heart J*. Jul 2012; 33(14):1787-1847.
87. Mehra MR, Naka Y, Uriel N, et al. A fully magnetically levitated circulatory pump for advanced heart failure. *N Engl J Med*. Feb 02 2017; 376(5):440-450.

88. Miller LW, Pagani FD, Russell SD et al. Use of a continuous-flow device in patients awaiting heart transplantation. *N Engl J Med* 2007; 357(9):885-896.
89. Mohamedali B, Bhat G, Yost G, et al. Survival on biventricular mechanical support with the Centrimag(R) as a bridge to decision: a single-center risk stratification. *Perfusion*. Apr 2015; 30(3):201-208.
90. Nativi JN, Drakos SG, Kucheryavaya AY et al. Changing outcomes in patients bridged to heart transplantation with continuous- versus pulsatile-flow ventricular assist devices: an analysis of the registry of the International Society for Heart and Lung Transplantation. *J Heart Lung Transplant* 2011; 30(8):854-61.
91. Netuka I, Sood P, Pya Y et al. Fully magnetically levitated left ventricular assist system for treating advanced HF: a multicenter study. *J Am Coll Cardiol*. Dec 15 2015; 66(23):2579-2589.
92. O'Neill WW, Kleiman NS, Moses J, et al. A prospective, randomized clinical trial of hemodynamic support with Impella 2.5 versus intra-aortic balloon pump in patients undergoing high-risk percutaneous coronary intervention: the PROTECT II study. *Circulation*. Oct 2 2012; 126(14):1717-1727.
93. O'Neill WW, Schreiber T, Wohns DH, et al. The current use of Impella 2.5 in acute myocardial infarction complicated by cardiogenic shock: results from the USpella Registry. *J Interv Cardiol*. Feb 2014; 27(1):1-11.
94. Organ Procurement and Transplantation Network. Heart Kaplan-Meier Patient Survival Rates For Transplants Performed : 2008 - 2015. 2018; <https://optn.transplant.hrsa.gov/data/view-data-reports/national-data/#>. Accessed August 7, 2018.
95. Ouweneel DM, Eriksen E, Sjauw KD, et al. Percutaneous mechanical circulatory support versus intra-aortic balloon pump in cardiogenic shock after acute myocardial infarction. *J Am Coll Cardiol*. Jan 24 2017; 69(3):278-287.
96. Park SJ, Tector A, Piccioni W et al. Left ventricular assist devices as destination therapy: a new look at survival. *J Thorac Cardiovasc Surg* 2005; 129(1):9-17.
97. Patel NJ, Singh V, Patel SV et al. Percutaneous coronary interventions and hemodynamic support in the USA: A 5 year experience. *J Interv Cardiol*. 2015 Dec; 28(6):563-73.
98. Patel ND, Weiss ES, Schaffer J et al. Right heart dysfunction after left ventricular assist device implantation: a comparison of the pulsatile HeartMate I and axial-flow HeartMate II devices. *Ann Thorac Surg* 2008; 86(3):832-40; discussion 32-40.
99. Peura JL, Colvin-Adams M, Francis GS, et al. Recommendations for the use of mechanical circulatory support: device strategies and patient selection: a scientific statement from the American Heart Association. *Circulation*. Nov 27 2012;126(22):2648-2667.
100. Poire, VL. The HeartMate left ventricular assist system: Worldwide clinical results. *European Journal of Cardio-Thoracic Surgery* 1997; 11 Suppl: S39-S44.
101. Ponikowski P, Voors AA, Anker SD, et al. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC). Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur Heart J*. Jul 14 2016; 37(27):2129-2200.
102. Pruijsten RV, Lok SI, Kirkels HH et al. Functional and haemodynamic recovery after implantation of continuous-flow left ventricular assist devices in comparison with

- pulsatile left ventricular assist devices in patients with end-stage heart failure. *Eur J Heart Fail* 2012; 14(3):319-25.
103. Reddy YM, Chinitz L, Mansour M, et al. Percutaneous left ventricular assist devices in ventricular tachycardia ablation: multicenter experience. *Circ Arrhythm Electrophysiol.* Apr 2014;7(2):244-250.
 104. Rihal CS, Naidu SS, Givertz MM, et al. 2015 SCAI/ACC/HFSA/STS Clinical Expert Consensus Statement on the Use of Percutaneous Mechanical Circulatory Support Devices in Cardiovascular Care: Endorsed by the American Heart Association, the Cardiological Society of India, and Sociedad Latino Americana de Cardiologia Intervencion; Affirmation of Value by the Canadian Association of Interventional Cardiology-Association Canadienne de Cardiologie d'intervention. *J Am Coll Cardiol.* May 19 2015; 65(19):e7-e26.
 105. Rogers JG, Butler J, Lansman SL, et al. Chronic mechanical circulatory support for Inotrope-dependent heart failure patients who are not transplant candidates- Results of the INTREPID Trial. *J Am Coll Cardiol.* 2007; 50(8):741-747.
 106. Rogers JG, Pagani FD, Tatroles AJ, et al. Intrapericardial left ventricular assist device for advanced heart failure. *N Engl J Med.* Feb 02 2017; 376(5):451-460.
 107. Romeo F, Acconcia MC, Sergi D, et al. Percutaneous assist devices in acute myocardial infarction with cardiogenic shock: Review, meta-analysis. *World J Cardiol.* Jan 26 2016; 8(1):98-111.
 108. Rose EA, Gelijns AC, Moskowitz AJ, et al. Long-term use of a left ventricular assist device for end-state heart failure. *The New England Journal of Medicine*, November 2001, Vol. 345, No. 20: 1435-43.
 109. Schafer A, Werner N, Burkhoff D, et al. Influence of Timing and Predicted Risk on Mortality in Impella-Treated Infarct-Related Cardiogenic Shock Patients. *Front Cardiovasc Med.* 2020; 7: 74.
 110. Schreiber T, Wah Htun W, Blank N, et al. Real-world supported unprotected left main percutaneous coronary intervention with impella device; data from the USpella Registry. *Catheter Cardiovasc Interv.* Apr 18 2017; 90(4):576-581.
 111. Seyfarth M, Sibbing D, Bauer I, et al. A randomized clinical trial to evaluate the safety and efficacy of a percutaneous left ventricular assist device versus intra-aortic balloon pumping for treatment of cardiogenic shock caused by myocardial infarction. *J Am Coll Cardiol.* Nov 4 2008; 52(19):1584-1588.
 112. Shuhaiber JH, Hur K, Gibbons R. The influence of preoperative use of ventricular assist devices on survival after heart transplantation: propensity score matched analysis. *BMJ* 2010; 340:c392.
 113. Shuhaiber JH, Jenkins D, Berman M et al. The Papworth experience with the Levitronix CentriMag ventricular assist device. *J Heart Lung Transplant* 2008; 27(2):158-64.
 114. Sieweke JT, Berliner D, Tongers J, et al. Mortality in patients with cardiogenic shock treated with the Impella CP microaxial pump for isolated left ventricular failure. *Eur Heart J Acute Cardiovasc Care.* Mar 2020; 9(2): 138-148.
 115. Sjaww KD, Konorza T, Erbel R, et al. Supported high-risk percutaneous coronary intervention with the Impella 2.5 device the Europella registry. *J Am Coll Cardiol.* Dec 15 2009; 54(25):2430-2434.

116. Slaughter MS, Pagani FD, McGee EC, et al. HeartWare ventricular assist system for bridge to transplant: combined results of the bridge to transplant and continued access protocol trial. *J Heart Lung Transplant*. Jul 2013; 32(7):675-683.
117. Slaughter MS, Rogers JG, Milano CA et al. Advanced heart failure treated with continuous-flow left ventricular assist device. *N Engl J Med* 2009; 361(23):2241-51.
118. Starling RC, Estep JD, Horstmanshof DA, et al. Risk Assessment and comparative effectiveness of left ventricular assist device and medical management in ambulatory heart failure patients: The ROADMAP Study 2-year results. *JACC Heart Fail*. Jul 2017; 5(7):518-527.
119. Starling RC, Naka Y, Boyle AJ, et al. Results of the Post-U.S. Food and Drug Administration-Approval Study With a Continuous Flow Left Ventricular Assist Device as a Bridge to Heart Transplantation: A Prospective Study Using the INTERMACS (Interagency Registry for Mechanically Assisted Circulatory Support). *Journal of the American College of Cardiology*. 2011; 57(19):1890-1898.
120. Struber M, Sander K, Lahpor J et al. HeartMate II left ventricular assist device; early European experience. *Eur J Cardiothorac Surg* 2008; 34(2):289-94.
121. Strueber M, O'Driscoll G, Jansz P et al. Multicenter evaluation of an intrapericardial left ventricular assist system. *J Am Coll Cardiol* 2011; 57(12):1375-82.
122. Takayama H, Soni L, Kalesan B, et al. Bridge-to-decision therapy with a continuous-flow external ventricular assist device in refractory cardiogenic shock of various causes. *Circ Heart Fail*. Sep 2014; 7(5):799-806.
123. TEC Assessment Program. Left ventricular assist devices as destination therapy for end-stage heart failure. 2002;Volume 17;Tab 19.
124. TEC Assessment Program. Ventricular assist devices in bridging to heart transplantation. 1996;Volume 11;Tab 26.
125. Thiele H, Sick P, Boudriot E, et al. Randomized comparison of intra-aortic balloon support with a percutaneous left ventricular assist device in patients with revascularized acute myocardial infarction complicated by cardiogenic shock. *Eur Heart J*. Jul 2005; 26(13):1276-1283.
126. Topkara VK, Garan AR, Fine B, et al. Myocardial recovery in patients receiving contemporary left ventricular assist devices: results from the Interagency Registry for Mechanically Assisted Circulatory Support (INTERMACS). *Circ Heart Fail*. Jul 2016;9(7).
127. Torregrossa G, Morshuis M, Varghese R, et al. Results with SynCardia total artificial heart beyond 1 year. *ASAIO J*. Nov-Dec 2014; 60(6):626-634.
128. U.S. Food and Drug Administration (FDA). Impella Recover LP 2.5 Percutaneous Cardiac Support System 510(k) approval. www.fda.gov. Last reviewed March 2012.
129. Ventura PA, Alharethi R, Budge D et al. Differential impact on post-transplant outcomes between pulsatile- and continuous-flow left ventricular assist devices. *Clin Transplant* 2011; 25(4):E390-5.
130. Wehman B, Stafford KA, Bittle GJ, et al. Modern outcomes of mechanical circulatory support as a bridge to pediatric heart transplantation. *Ann Thorac Surg*. Jun 2016; 101(6):2321-2327.
131. Wever-Pinzon O, Drakos SG, McKellar SH, et al. Cardiac recovery during long-term left ventricular assist device support. *J Am Coll Cardiol*. Oct 04 2016; 68(14):1540-1553.

132. Wieselthaler GM, Schima H, Lassnigg AM et al. Lessons learned from the first clinical implants of the DeBakey ventricular assist device axial pump: a single center report. *Ann Thorac Surg* 2001; 71(3 Suppl):S139-43; discussion S44-6.
133. Yancy CW, Jessup M, Bozkurt B, et al. 2013 ACCF/AHA Guideline for the Management of Heart Failure-A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2013; 62(16):e147-e239.

POLICY HISTORY:

Adopted for Blue Advantage, February 2016

Available for comment February 15 through March 30, 2016

Medical Policy Group, August 2016

Medical Policy Group, September 2017

Medical Policy Group, February 2018

Medical Policy Group, September 2020

Medical Policy Group, November 2020: Annual Coding Update. Added new CPT codes 33995 and 33997. Revised CPT codes 33990-33993 to clarify left or right heart.

Medical Policy Group, April 2021

Medical Policy Group, August 2021

Medical Policy Group, December 2021: 2022 Annual Coding Update. Moved CPT codes from Current coding section. Updated Previous Coding section to include codes 0451T, 0452T, 0453T, 0454T, 0455T, 0456T, 0457T, 0458T, 0459T, 0460T, 0461T, and 0462T. Added unlisted CPT Code 33999 to Current Coding section. CPT Code 33999 will be used to report permanently implantable aortic counterpulsation ventricular assist devices after 1/1/2022.

This medical policy is not an authorization, certification, explanation of benefits, or a contract. Eligibility and benefits are determined on a case-by-case basis according to the terms of the member's plan in effect as of the date services are rendered. All medical policies are based on (i) research of current medical literature and (ii) review of common medical practices in the treatment and diagnosis of disease as of the date hereof. Physicians and other providers are solely responsible for all aspects of medical care and treatment, including the type, quality, and levels of care and treatment.

This policy is intended to be used for adjudication of claims (including pre-admission certification, pre-determinations, and pre-procedure review) in Blue Cross and Blue Shield's administration of plan contracts.