



BlueCross BlueShield
of Alabama

Name of Blue Advantage Policy:

Transcatheter Closure Devices for Septal Defects

Policy #: 218
Category: Medical

Latest Review Date: June 2020
Policy Grade: B

BACKGROUND:

Blue Advantage medical policy does not conflict with Local Coverage Determinations (LCDs), Local Medical Review Policies (LMRPs) or National Coverage Determinations (NCDs) or with coverage provisions in Medicare manuals, instructions or operational policy letters. In order to be covered by Blue Advantage the service shall be reasonable and necessary under Title XVIII of the Social Security Act, Section 1862(a)(1)(A). The service is considered reasonable and necessary if it is determined that the service is:

1. *Safe and effective;*
2. *Not experimental or investigational*;*
3. *Appropriate, including duration and frequency that is considered appropriate for the service, in terms of whether it is:*
 - *Furnished in accordance with accepted standards of medical practice for the diagnosis or treatment of the patient's condition or to improve the function of a malformed body member;*
 - *Furnished in a setting appropriate to the patient's medical needs and condition;*
 - *Ordered and furnished by qualified personnel;*
 - *One that meets, but does not exceed, the patient's medical need; and*
 - *At least as beneficial as an existing and available medically appropriate alternative.*

Routine costs of qualifying clinical trial services with dates of service on or after September 19, 2000 which meet the requirements of the Clinical Trials NCD are considered reasonable and necessary by Medicare. Providers should bill **Original Medicare for covered services that are related to **clinical trials** that meet Medicare requirements (Refer to Medicare National Coverage Determinations Manual, Chapter 1, Section 310 and Medicare Claims Processing Manual Chapter 32, Sections 69.0-69.11).*

POLICY:

Effective for dates of service on or after August 14, 2016:

Blue Advantage will treat **transcatheter closure of a patent foramen ovale (PFO) using an FDA approved closure device** as a **covered benefit** for the following indication:

- Diagnosis of PFO with a right to left shunt confirmed by echocardiography
- Diagnosis of cryptogenic stroke due to a presumed paradoxical embolism

Blue Advantage will treat **transcatheter closure of atrial septal defects (ASD) in adult or pediatric patients** as a **covered benefit** for any of the following indications:

- For the occlusion of ASDs in secundum position OR patients who have undergone a fenestrated Fontan procedure who now require closure of the fenestration;
- Have echocardiographic evidence of ostium secundum ASD AND clinical evidence of right ventricular volume overload (e.g. 1.5:1 degree of left to right shunt or RV enlargement).
- Have echocardiographic evidence of ostium secundum ASD AND clinical evidence of paradoxical embolism
- Documented platypnea- orthodeoxia
- Presence of net left-to-right shunting, pulmonary artery pressure less than two thirds systemic levels, PVR less than two thirds systemic vascular resistance, or when responsive to either pulmonary vasodilator therapy or test occlusion of the defect

Blue Advantage will treat **transcatheter closure of atrial septal defects (ASD,) in all other situations, including but not limited to coronary sinus ASD, ostium primum ASD and sinus venosus ASD,** as a **non-covered benefit** and as **investigational**.

Blue Advantage will treat **transcatheter closure of ventricular septal defects (VSD)** as a **covered benefit** for any of the following indications:

- Complex VSD of significant size to warrant closure (e.g. large volume left to right shunt, pulmonary hypertension, and/or clinical symptoms of congestive heart failure) **AND** are considered to be at high risk for standard transatrial or transarterial surgical closure based on anatomical conditions and/or based on overall medical condition.**
- Iatrogenic artifacts after surgical replacement of the aortic valve

**High risk anatomical factors for transatrial or transarterial surgical closure include patients:

- Requiring left ventriculotomy or an extensive right ventriculotomy
- With a failed previous VSD closure
- With multiple apical and/or anterior muscular VSDs (Swiss Cheese Septum)
- With posterior apical VSDs covered by trabeculae

Blue Advantage will treat **transcatheter closure of ventricular septal defects (VSD)** as a **non-covered benefit** and as **investigational** including but not limited to, when the above criteria are not met and in patients with severe irreversible PAH.

Blue Advantage will treat **transcatheter closure of patent ductus arteriosus (PDA)** as a **covered benefit** when using an FDA approved device for this indication.

Blue Advantage does not approve or deny procedures, services, testing, or equipment for our members. Our decisions concern coverage only. The decision of whether or not to have a certain test, treatment or procedure is one made between the physician and his/her patient. Blue Advantage administers benefits based on the members' contract and medical policies. Physicians should always exercise their best medical judgment in providing the care they feel is most appropriate for their patients. Needed care should not be delayed or refused because of a coverage determination.

DESCRIPTION OF PROCEDURE OR SERVICE:

Despite the success of standard open-heart surgery to repair cardiac defects, the risks and morbidity of open-heart surgery remain. Interventional cardiac catheterization techniques have advanced to a point where percutaneous transcatheter devices can be offered as an alternative for carefully selected patients. Transcatheter closure devices are permanent implants designed to close defects between chambers of the heart, including atrial septal defect (ASD), ventricular septal defect (VSD), and patent foramen ovale (PFO). Devices are also used for persistent patent ductus arteriosus (PDA), which is an opening between the descending thoracic aorta and the pulmonary artery which fails close after birth. These devices are self-expandable, self-centering umbrella-like devices. They are implanted in the defect through catheters inserted into either a vein or an artery using a transcatheter or percutaneous approach.

The standard for managing clinically significant cardiac defects mentioned above has been surgical closure, which except for complex ventricular septal defects, is associated with a very low mortality. Conventional surgical closure is done through a midline sternotomy. The newer approaches utilizing the closure devices offer repair of the defect without major thoracic surgery, less post-operative pain, and decreased hospital stay without compromising outcomes in a select group of patients.

Patent Foramen Ovale (PFO)

The foramen ovale, a component of fetal cardiovascular circulation, consists of a communication between the right and left atrium that functions as a vascular bypass of the un-inflated lungs. The ductus arteriosus is another feature of the fetal cardiovascular circulation, consisting of a connection between the pulmonary artery and the distal aorta. Prior to birth, the foramen ovale is held open by the large flow of blood into the left atrium from the inferior vena cava. Over a course of months after birth, an increase in left arterial pressure and a decrease in right atrial pressure result in the permanent closure of the foramen ovale in most individuals. However, a patent foramen ovale may be detected in up to 25% of asymptomatic adults. In some epidemiologic studies, patent foramen ovale (PFO) has been associated with cryptogenic stroke, a type of stroke defined as an ischemic stroke occurring in the absence of potential cardiac, pulmonary, vascular, or neurologic sources. Studies also show an association of PFO and migraine headache.

Treatment

Conventional therapy for cryptogenic stroke consists of antiplatelet therapy (aspirin, clopidogrel, or dipyridamole given alone or in combination) or oral anticoagulation with warfarin. In general, patients with a known clotting disorder or evidence of preexisting thromboembolism are treated with warfarin, and patients without these risk factors are treated with antiplatelet agents. Closure devices are nonpharmacologic alternatives to medical therapy for cryptogenic stroke in patients with a PFO.

Atrial Septal Defect (ASD)

Unlike PFOs, which represents the postnatal persistence of normal fetal cardiovascular physiology, ASDs represent an abnormality in the development of the heart that results in free communication between the atria. ASDs are categorized according to their anatomy. Ostium secundum describes defects that are located mid-septally and are typically near the fossa ovalis. Ostium primum defects lie immediately adjacent to the atrioventricular valves and are within the spectrum of atrioventricular septal defects. Primum defects occur commonly in patients with Down syndrome. Sinus venous defects occur high in the atrial septum and are frequently associated with anomalies of the pulmonary veins.

Ostium secundum ASDs are the third most common form of congenital heart disorder and one of the most common congenital cardiac malformations in adults, accounting for 30% to 40% of these patients older than age 40 years. The ASD often goes unnoticed for decades because the physical signs are subtle and the clinical sequelae are mild. However, virtually all patients who survive into their sixth decade are symptomatic; fewer than 50% of patients survive beyond age 40 to 50 years due to heart failure or pulmonary hypertension related to the left-to-right shunt. Symptoms related to ASD depend on the size of the defect and the relative diastolic filling properties of the left and right ventricles. Reduced left ventricular compliance and mitral stenosis will increase left-to-right shunting across the defect. Conditions that reduce right ventricular compliance and tricuspid stenosis will reduce left-to-right shunting or cause a right-to-left shunt. Symptoms of an ASD include exercise intolerance and dyspnea, atrial fibrillation, and, less commonly, signs of right heart failure. Patients with ASDs are also at risk for paradoxical emboli.

Treatment

Repair of ASDs is recommended for those with a pulmonary to systemic flow ratio ($Q_p:Q_s$) exceeding 1.5:1.0. Despite the success of operative repair, there has been interest in developing a catheter-based approach to ASD repair to avoid the risks and morbidity of open heart surgery. A variety of devices have been researched. Technical challenges include minimizing the size of device so that smaller catheters can be used, developing techniques to properly center the device across the ASD, and ensuring that the device can be easily retrieved or repositioned, if necessary.

Individuals with ASDs and a history of cryptogenic stroke are typically treated with antiplatelet agents, given an absence of evidence that systemic anticoagulation is associated with outcome improvements.

Transcatheter Closure Devices

Transcatheter PFO and ASD occluders consist of a single or paired wire mesh discs that are covered or filled with polyester or polymer fabric that are placed over the septal defect. Over

time, the occlusion system is epithelialized. ASD occluder devices consist of flexible mesh disks that are delivered via catheter to cover the ASD.

Ventricular Septal Defect (VSD)

VSD is the most common congenital heart defect at birth and presents in approximately 42 out of every 10,000 births per year. Although VSD is most often an isolated lesion, it is a common component of complex abnormalities such as conotruncal defects (e.g., tetralogy of Fallot and Transposition of Great Arteries). VSD can also be associated with left-sided obstructive lesions such as subaortic stenosis (SubAS) and coarctation of the aorta. A subpulmonary (supracristal) VSD is often associated with progressive aortic valve regurgitation caused by prolapse of the aortic cusp (usually right) through the defect.

Ventricular septal defects (VSDs) are usually present at birth but may also occur following myocardial infarction. Small VSDs may never be detected. A large VSD can allow blood to flow from the left to the right ventricle which increases load on the heart and lungs. Septal defects may also be created during surgical repair of certain types of congenital heart defects, such as hypoplastic left ventricle syndrome. A hole is made in the septum of the repaired section of the heart (Fontan fenestration) to improve recovery from the surgery. The fenestration is intended to be closed at a later time.

Treatment

Small VSDs may cause no problems and close on its own. If the VSD is large, open heart surgery may be required which is usually performed in infancy or childhood. During open heart surgery, a patch is sewn over the defect or if possible, the defect is sewn shut without a patch.

Transcatheter Closure Devices

The Amplatzer™ Muscular VSD Occluder is a device to occlude ventricular septal defects. The device is made of a polyester material that promotes occlusion and tissue in-growth. Similarly, the CardioSEAL® Septal Occlusion System is a permanent implant made of a metal framework to occlude VSDs.

Patent Ductal Arteriosus (PDA)

PDA is another fairly common type of congenital heart defect (CHD) affecting approximately 8 out of every 1000 premature babies. Although it can affect full term babies, it is more common in premature babies. With this CHD, there is abnormal blood flow between the aorta and pulmonary artery. These two arteries are connected by the ductus arteriosus which is a normal and essential part of fetal blood circulation. Soon after birth, the vessel should close. When the vessel does not close, it allows oxygen rich blood from the aorta to mix with oxygen poor blood from the pulmonary artery which may strain the heart and increase blood pressure in lung arteries.

Treatment

A PDA may be treated with medications, catheter based procedures or open surgery. Transcatheter closure of a PDA is an established treatment and is often the treatment of choice.

KEY POINTS:

The most recent update with literature review is through March 9, 2020.

Summary of Evidence

PFO

For individuals who have PFO and cryptogenic stroke who receive PFO closure with a transcatheter device, the evidence includes multiple randomized controlled trials (RCTs) comparing device-based PFO closure with medical therapy, systematic reviews and meta-analyses of these studies. Relevant outcomes include overall survival, morbid events, and treatment-related morbidity and mortality. The RCTs comparing PFO closure with medical management have suggested that PFO closure is more effective than medical therapy in reducing event rates. While these results were not statistically significant by intention-to-treat analyses in the first 3 trials (i.e., CLOSURE I, PC, and RESPECT [initial study]), they were statistically significant in later trials (i.e., RESPECT [extended follow-up], REDUCE, and CLOSE). Use of appropriate patient selection criteria to eliminate other causes of cryptogenic stroke in RESPECT, REDUCE, and CLOSE trials contributed to findings of the superiority of PFO closure compared with medical management. Of note, higher rates of atrial fibrillation were reported in a few of the individual trials and in the meta-analysis that incorporated evidence from RESPECT, REDUCE, and CLOSE trials. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have PFO and migraines who receive PFO closure with a transcatheter device, the evidence includes randomized controlled trials of PFO closure, along with multiple observational studies reporting on the association between PFO and migraine. Relevant outcomes are symptoms, quality of life, medication use, and treatment-related morbidity and mortality. The available sham-controlled RCT did not demonstrate significant improvements in migraine symptoms after PFO closure. A second RCT with blinded endpoint evaluation did not demonstrate improvements in migraine days after PFO closure, but was likely underpowered. Nonrandomized studies have shown highly variable rates of migraine improvement after PFO closure. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have PFO and conditions associated with PFO other than cryptogenic stroke or migraine (e.g., myocardial infarction with normal coronary arteries, decompression illness, high altitude pulmonary edema, obstructive sleep apnea) who receive PFO closure with a transcatheter device, the evidence includes small case series and case reports. Relevant outcomes are symptoms, change in disease status, morbid events, and treatment-related morbidity and mortality. The body of evidence consists only of small case series and case reports. Comparative studies are needed to evaluate outcomes in similar patient groups treated with and without PFO closure. The evidence is insufficient to determine the effects of the technology on health outcomes.

ASD

For individuals who have ASD and evidence of left-to-right shunt or right-ventricular overload who receive ASD closure with a transcatheter device, the evidence includes nonrandomized comparative studies and single-arm studies. Relevant outcomes are symptoms, change in

disease status, and treatment-related morbidity and mortality. The available nonrandomized comparative studies and single-arm case series show high success rates of closure using closure devices approaching the high success rates of surgery, which are supported by meta-analyses of these studies. The percutaneous approach has a low complication rate and avoids the morbidity and complications of open surgery. If the percutaneous approach is unsuccessful, ASD closure can be achieved using surgery. Because of the benefits of percutaneous closure over open surgery, it can be determined that transcatheter ASD closure improves outcomes in patients with an indication for ASD closure. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

VSD

The evidence for transcatheter closure of VSD has shown high closure rates, low procedural mortality and positive short term results. Low complication rates and shorter hospital stays have made this procedure more favorable over the surgical treatment when appropriate. The evidence is sufficient to determine qualitatively that the technology results in a meaningful improvement in the net health outcome.

PDA

The evidence for transcatheter occlusion of PDA has shown that this is a well established treatment, and is an efficient and safe procedure. Studies have shown excellent results for short and long term studies. Low complication rates and shorter hospital stays have made this procedure more favorable over the surgical treatment when appropriate. The evidence is sufficient to determine qualitatively that the technology results in a meaningful improvement in the net health outcome.

Practice Guidelines and Position Statements

PFO

American College of Chest Physicians

In 2012, the American College of Chest Physicians updated its guidelines on antithrombotic therapy and the prevention of thrombosis, which made the following recommendations related to patent foramen ovale (PFO) and cryptogenic stroke:

- “We suggest that patients with stroke and PFO are treated with antiplatelet therapy following the recommendations for patients with noncardioembolic stroke.... In patients with a history of noncardioembolic ischemic stroke or TIA, we recommend long-term treatment with aspirin (75-100 mg once daily), clopidogrel (75 mg once daily), aspirin/extended release dipyridamole (25 mg/200 mg bid), or cilostazol (100 mg bid) over no antiplatelet therapy (Grade 1A), oral anticoagulants (Grade 1B), the combination of clopidogrel plus aspirin (Grade 1B), or triflusal (Grade 2B).”

American Academy of Neurology

In 2016, the American Academy of Neurology (AAN) issued updated evidence-based guidelines about the management of patients with stroke and PFO to address whether percutaneous closure of PFO is superior to medical therapy alone. Following a systematic review of the literature and structured formulation of recommendations, the AAN’s developed

conclusions for the Amplatzer PFO Occluder devices. For patients with cryptogenic stroke and PFO, percutaneous PFO closure with the Amplatzer PFO Occluder:

- “Possibly decreases the risk of recurrent stroke—RD -1.68%, 95% CI -3.18% to -0.19%;
- “Possibly increases the risk of new-onset AF—RD 1.64%, 95% CI 0.07%–3.2% (2 Class I studies; confidence downgraded to low for risk of bias relative to magnitude of effect and imprecision);”
- “Is highly likely to be associated with a procedural complication risk of 3.4%, 95% CI 2.3%–5% (2 Class I studies).”

The guidelines conclude:

“Clinicians should not routinely offer percutaneous PFO closure to patients with cryptogenic ischemic stroke outside of a research setting (Level R). In rare circumstances, such as recurrent strokes despite adequate medical therapy with no other mechanism identified, clinicians may offer the AMPLATZER PFO Occluder if it is available (Level C).”

American Heart Association and American Stroke Association

In 2014, the American Heart Association (AHA) and American Stroke Association published updated guidelines on the prevention of stroke in patients with ischemic stroke or TIA. The guidelines list the following recommendations for device-based closure for patent foramen ovale (PFO):

- “For patients with a cryptogenic ischemic stroke or TIA and a PFO without evidence for DVT, available data do not support a benefit for PFO closure (Class III; Level of Evidence A).”
- “In the setting of PFO and DVT, PFO closure by a transcatheter device might be considered, depending on the risk of recurrent DVT (Class IIb; Level of Evidence C).”

ASD

American College of Cardiology and American Heart Association

In 2018, the AHA and ACC published guidelines for the management of adults with congenital heart disease. They recommended:

American College of Cardiology and American Heart Association Recommendations for Treating Atrial Septal Defect

Condition	Recommendation	CORa/LOEb
Symptomatic isolated secundum ASD, right atrial and/or RV enlargement, and net left-to-right shunt sufficiency large enough to cause physiological sequelae, without cyanosis at rest or during exercise	Transcatheter or surgical closure	I1/B-NR2

Symptomatic primum ASD, sinus venosus defect, or coronary sinus defect, right atrial and/or RV enlargement, and net left-to-right shunt sufficiency large enough to cause physiological sequelae, without cyanosis at rest or during exercise	Surgical closure unless precluded by comorbidities	I1/B-NR2
Asymptomatic isolated secundum ASD, right atrial and RV enlargement, and net left-to-right shunt sufficiency large enough to cause physiological sequelae, without cyanosis at rest or during exercise	Transcatheter or surgical closure	I1a1/C-LD2
Secundum ASD when a concomitant surgical procedure is being performed and there is a net left-to-right shunt sufficiently large enough to cause physiological sequelae, and right atrial and RV enlargement without cyanosis at rest or during exercise	Surgical closure	I1a1/C-LD2
ASD when net left-to-right shunt is $\geq 1.5:1$, PA systolic pressure and/or pulmonary vascular resistance is greater than of one-third of systemic resistance	Percutaneous or surgical closure	I1b1/B-NR2
ASD with PA systolic pressure greater than two-thirds systemic, pulmonary vascular resistance greater than two-thirds systemic, and/or a net left-to-right shunt	ASD closure should not be performed	III-Harm1/C-LD2

Adapted from Stout et al (2019)

ASD: atrial septal defect; COR: class (strength) of recommendation; LOE: level (quality) of evidence; PA: pulmonary artery; RCT: randomized controlled trial; RV: right ventricular.

a COR key: I=strong; I1a=moderate; I1b=weak; III: No Benefit=weak; III: Harm=strong. b LOE key: A=high quality from >1 RCT, meta-analyses of high-quality RCTs, ≥ 1 RCT corroborated by high-quality registry studies; B-R=randomized, moderate-quality evidence from ≥ 1 RCT or meta-analysis of moderate-quality RCTs; B-NR=nonrandomized, moderate-quality evidence from ≥ 1 well-designed, well-executed nonrandomized study, observational study, or registry study, or meta-analyses of such studies; C-LD: limited data, randomized or nonrandomized observational or registry studies with limitations of design or execution, meta-analyses of such studies, or physiological or mechanistic studies in human subjects; C-EO: expert opinion

VSD

National Institute for Health and Care Excellence

In 2010, NICE issued a guideline that stated, “Current evidence on the safety and efficacy of transcatheter endovascular closure of perimembranous ventricular septal defect (VSD) is adequate to support the use of this procedure provided that normal arrangements are in place for clinical governance, consent and audit.”

American College of Cardiology and American Heart Association

In 2018, the AHA and ACC published guidelines for the management of adults with congenital heart disease. They did not have a specific recommendation for transcatheter VSD closure, but stated that “Transcatheter device occlusion of muscular and perimembranous VSD is feasible, and trials have demonstrated a good safety and efficacy profile.

PDA

In 2018, the AHA and ACC published guidelines for the management of adults with congenital heart disease. They did not have a specific recommendation for transcatheter PDA closure, but stated that:

“When signs of volume overload are indicative of significant left-to-right shunt, closing the PDA is likely to prevent further left atrial or LV enlargement, progression or development of PAH, and pulmonary hypertension secondary to left HF and will possibly provide symptom relief if symptoms are present. Closure is typically performed percutaneously with good success and minimal complications.”

The ACC/AHA issued guidance in 2008 regarding closure of PDA.

“Closure of a PDA either percutaneously or surgically is indicated for the following:

- a. Left atrial and/or LV enlargement or is PAH is present, or in the presence of net left to right shunting. (Level of Evidence C)
- b. Prior endocarditis (Level of Evidence C)

U.S. Preventive Services Task Force Recommendations

Use of closure devices are not a preventive service.

KEY WORDS:

Atrial Septal Defect, ASD, Ventricular Septal Defect, VSD, Patent Foramen Ovale, PFO, Patent Ductus Arteriosus, PDA, AMPLATZER Septal Occluder, Gore HELEX Septal Occluder, CardioSEAL Septal Occlusion System with Qwik Load, AMPLATZER Muscular VSO Occluder, CardioSEAL STAR Flex Septal Occlusion System, AMPLATZER PFO Occluder, Monodisc Occluder

APPROVED BY GOVERNING BODIES:

Patent Foramen Ovale

The U.S. Food and Drug Administration (FDA) has approved three devices for ASD closure through the premarket approval process or a premarket approval supplement: the Amplatzer Septal Occluder, the GORE HELEX Septal Occluder (discontinued), and the GORE CARDIOFORM Septal Occluder.

In March 2018, the FDA granted an expanded indication to the Gore® Cardioform Septal Occluder to reduce the risk of recurrent ischemic stroke. It is a permanently implanted device indicated for the percutaneous, transcatheter closure of the following defects of the atrial septum:

- Patent foramen ovale to reduce the risk of recurrent ischemic stroke in patients, predominately between the ages of 18 and 60 years, who have had a cryptogenic stroke due to a presumed paradoxical embolism, as determined by a neurologist and cardiologist following an evaluation to exclude known causes of ischemic stroke.

In November 2016, the FDA approved the Amplatzer PFO Occluder (Abbott) through the premarket approval (PMA) process for the following indication:

“For percutaneous transcatheter closure of a patent foramen ovale (PFO) to reduce the risk of recurrent ischemic stroke in patients, predominantly between the ages of 18 and 60 years, who have had a cryptogenic stroke due to a presumed paradoxical embolism, as determined by a neurologist and cardiologist following an evaluation to exclude known causes of ischemic stroke.”

In 2002, 2 transcatheter devices were cleared for marketing by the U.S. Food and Drug Administration (FDA) through a humanitarian device exemption as treatment for patients with cryptogenic stroke and PFO: the CardioSEAL® Septal Occlusion System (NMT Medical; device no longer commercially available) and the Amplatzer® PFO Occluder (Amplatzer, now Abbott). Following the limited FDA approval, use of PFO closure devices increased by more than 50-fold, well in excess of the 4000 per year threshold intended under the humanitarian device exemption, prompting FDA to withdraw the humanitarian device exemption approval for these devices in 2007.

Atrial Septal Defect

Three devices have been approved by the FDA through the premarket approval process or a premarket approval supplement for transcatheter ASD closure (see Table 1) (FDA product code: MLV).

Table 1. ASD Closure Devices Approved by the Food and Drug Administration

Device	Manufacturer	PMA Approval Date	Indications
Amplatzer™ Septal Occluder	St. Jude Medical (Abbott Medical)	Dec 2001	<ul style="list-style-type: none"> • Occlusion of ASDs in the secundum position • Use in patients who have had a fenestrated Fontan procedure who require closure of the fenestration • Patients indicated for ASD closure have echocardiographic evidence of ostium secundum ASD and clinical evidence of right ventricular volume overload.
GORE HELEX Septal Occluder	W.L. Gore & Associates	Aug 2006 (discontinued)	<ul style="list-style-type: none"> • Percutaneous, transcatheter closure of ostium secundum ASDs
GORE CARDIOFORM ASD Occluder (formerly GORE CARDIOFORM Septal Occluder)	W.L. Gore & Associates	May 2019 (supplement; name change) Oct 2016 (supp.)	<ul style="list-style-type: none"> • Percutaneous, transcatheter closure of ostium secundum ASDs

ASD: atrial septal defect; PMA: premarket approval.

Ventricular Septal Defect

AMPLATZER® Muscular VSD Occluder received FDA approval via premarket application (PMA) on September 7, 2007 for closure of complex VSD of significant size to warrant closure on patients who are considered to be at high risk for standard transatrial or transarterial surgical closure based on anatomical conditions and/or based on overall medical condition.

CardioSEAL® Septal Occlusion System with Qwik Load (Nitinol Medical Technologies, Inc.) received FDA approval via premarket application (PMA) on December 5, 2001 for closure of complex VSD; however, NMT Medical, Inc ceased operations in 2011.

Patent Ductus Arteriosus

In May 2003, the Amplatzer Duct Occluder 180* Delivery System (AGA Medical Corporation, Golden Valley, MN) received FDA approval through the PMA process for the closure of PDA.

In August 2013, the Nit-Occlud® PDA (PFM medical) received FDA approval via PMA for closure of small to moderate size PDA

BENEFIT APPLICATION:

Coverage is subject to member’s specific benefits. Group specific policy will supersede this policy when applicable.

CURRENT CODING:

CPT Codes:

93799	Unlisted cardiovascular service or procedure
93580	Percutaneous transcatheter closure of congenital interatrial communication (i.e., Fontan fenestration, atrial septal defect) with implant
93581	Percutaneous transcatheter closure of a congenital ventricular septal defect with implant
93582	Percutaneous transcatheter closure of patent ductus arteriosus
93315	Transesophageal echocardiography for congenital cardiac anomalies; including probe placement, image acquisition, interpretation and report
93462	Left heart catheterization by transseptal puncture through intact septum or by transapical puncture (List separately in addition to code for primary procedure).
93563	Injection procedure during cardiac catheterization including imaging supervision, interpretation, and report; for selective coronary angiography during congenital heart catheterization (list separately in addition to code for primary procedure).
93568	Injection procedure during cardiac catheterization including imaging supervision, interpretation, and report; for pulmonary angiography (list separately in addition to code for primary procedure)

REFERENCES:

1. Abaci A, Unlu S, Alsancak Y, et al. Short and long term complications of device closure of atrial septal defect and patent foramen ovale: meta-analysis of 28,142 patients from 203 studies. *Catheter Cardiovasc Interv.* Dec 1 2013; 82(7):1123-1138.
2. Abbott. Amplatzer™ Muscular VSD Occluder. www.sjm.com/en/professionals/featured-products/structural-heart/structural-heart-occluders/vsd-closure-devices/amplatzer-muscular-vsd-occluder. Accessed June 15, 2018.
3. Albers GW, Amarenco P, Easton D, et al. Antithrombotic and thrombolytic therapy for ischemic stroke: American College of Chest physicians evidence-based clinical practice guidelines (8th edition). *Chest* 2008; 133; 630S-669S DOI 10.1378/chest.08-0720.
4. Alushi B, Biasco L, Orzan F, et al. Patent foramen ovale treatment strategy: an Italian large prospective study. *J Cardiovasc Med (Hagerstown)*. Jun 27 2014.
5. Alushi, BB, Lauten, AA, Cassese, SS, et al. Patent foramen ovale closure versus medical therapy for prevention of recurrent cryptogenic embolism: updated meta-analysis of randomized clinical trials. *Clin Res Cardiol*, 2018 Apr 13;107(9).
6. American College of Chest Physicians. Comparing outcomes of medical management and percutaneous closure in patients with patent foramen ovale and cryptogenic stroke. *Chest* 2008; journal.publications.chestnet.org/article.aspx?articleid=1094512&resultClick=3. Accessed July 2015.
7. American Heart Association. Ventricular Septal Defect. www.heart.org/HEARTORG/Conditions/CongenitalHeartDefects/AboutCongenitalHeartDefects/Ventricular-Septal-Defect-VSD_UCM_307041_Article.jsp#.WyPUiIVKhtQ. Accessed June 15, 2018.
8. Baruteau AE, Petit J, Lambert V, et al. Transcatheter closure of large atrial septal defects: feasibility and safety in a large adult and pediatric population. *Circ Cardiovasc Interv.* Dec 2014; 7(6):837-843.
9. Behjati-Ardakani M, Rafiei M, et al. Long-term results of transcatheter closure of patent ductus arteriosus in adolescents and adults with amplatzer duct occluder. *N Am J Med Sci.* 2015 May; 7(5): 208-11.
10. Berger F., Ewert P., et al. Percutaneous closure of large atrial septal defects with the Amplatzer septal occluder: technical overkill or recommendable alternative treatment? *J of Interventional Cardio*, 2001; 14:1.
11. Berger F, Vogel M, et al. Comparison of results and complications of surgical and Amplatzer device closure of atrial septal defects. *J Thorac Cardiovasc Surg* 1999; 118(4):674-8.
12. Biasco L, Infantino V, Orzan F, et al. Impact of transcatheter closure of patent foramen ovale in the evolution of migraine and role of residual shunt. *J Cardiol.* Apr 5 2014.
13. Brochu MC, Baril JF, et al. Improvement in exercise capacity in asymptomatic and mildly symptomatic adults after atrial septal defect percutaneous closure. *Circulation* 2002; 106(14):1821-6.
14. Butera G., Bini M.R., et al. Transcatheter closure of patent foramen ovale in patients with cryptogenic stroke, *Ital Heart J* 2001; 2 (2): 115-118.
15. Butera G, Biondi-Zoccai G, Sangiorgi G, et al. Percutaneous versus surgical closure of secundum atrial septal defects: a systematic review and meta-analysis of currently available clinical evidence. *EuroIntervention.* Jul 2011; 7(3):377-385.

16. Butera G, Saracino A, Danna P, et al. Transcatheter PFO closure with GORE ((R)) septal occluder: early and mid-term clinical results. *Catheter Cardiovasc Interv.* Nov 15 2013; 82(6):944-949.
17. Cao Q.L., Du Z.D., et al. Immediate and six-month results of the profile of the Amplatzer septal occluder as assessed by transesophageal echocardiography, *American Journal of Cardiology*, 2001; 88:1.
18. Capodanno D, Milazzo G, Vitale L, et al. Updating the evidence on patent foramen ovale closure versus medical therapy in patients with cryptogenic stroke: a systematic review and comprehensive meta-analysis of 2,303 patients from three randomised trials and 2,231 patients from 11 observational studies. *EuroIntervention.* Mar 20 2014; 9(11):1342-1349.
19. Carroll JD, Saver JL, Thaler DE et al. Closure of patent foramen ovale versus medical therapy after cryptogenic stroke. *N Engl J Med* 2013; 368(12):1092-100.
20. Centers for Disease Control and Prevention. Congenital heart defects (CHDs). Available at [//www.cdc.gov/ncbddd/heartdefects/ventricularseptaldefect.html](http://www.cdc.gov/ncbddd/heartdefects/ventricularseptaldefect.html). Accessed April 2016.
21. Chen TH, Hsiao YC, Cheng CC, et al. In-Hospital and 4-Year Clinical Outcomes Following Transcatheter Versus Surgical Closure for Secundum Atrial Septal Defect in Adults: A National Cohort Propensity Score Analysis. *Medicine (Baltimore)*. Sep 2015;94(38):e1524.
22. Chen L, Luo S, Yan L, et al. A systematic review of closure versus medical therapy for preventing recurrent stroke in patients with patent foramen ovale and cryptogenic stroke or transient ischemic attack. *J Neurol Sci.* Feb 15 2014; 337(1-2):3-7.
23. Cifarelli A, Musto C, Parma A, et al. Long-term outcome of transcatheter patent foramen ovale closure in patients with paradoxical embolism. *Int J Cardiol.* Jun 11 2010; 141(3):304-310.
24. Cotran R.S., Kumar V., and Collins T. *Robbins pathologic basis of disease*, W. B. Saunders Co. Philadelphia, 1999.
25. Demkow M., Ruzyllo W., et al. Transvenous closure of moderate and large secundum atrial septal defects in adults using the Amplatzer septal occluder, *Catheterization and Cardiovascular Interventions*, 2001; 52:188-193.
26. De Rosa S, Sievert H, Sabatino J, et al. Percutaneous closure versus medical treatment in stroke patients with patent foramen ovale: a systematic review and meta-analysis. *Ann Intern Med.* Mar 6 2018;168(5):343-350.
27. Dowson A, Mullen MJ, Peatfield R, et al. Migraine Intervention With STARFlex Technology (MIST) trial: a prospective, multicenter, double-blind, sham-controlled trial to evaluate the effectiveness of patent foramen ovale closure with STARFlex septal repair implant to resolve refractory migraine headache. *Circulation.* Mar 18 2008; 117(11):1397-1404.
28. Du ZD, Hijazi ZM, et al. Comparison between transcatheter and surgical closure of secundum atrial septal defect in children and adults: results of a multicenter nonrandomized trial. *J Am Coll Cardiol* 2002; 39(11):1836-44.
29. Du ZD, Koenig P, et al. Comparison of transcatheter closure of secundum atrial septal defect using the Amplatzer septal occluder associated with deficient versus sufficient rims. *Am J Cardiol* 2002; 90(8):865-9.
30. Eeckhout E, Martin S, Delabays A, et al. Very long-term follow-up after percutaneous closure of patent foramen ovale. *EuroIntervention.* Jan 17 2014.

31. Fischer G, Stieh J, et al. Experience with transcatheter closure of secundum atrial septal defects using the Amplatzer septal occluder: a single centre study in 236 consecutive patients. *Heart* 2003; 89(2):199-204.
32. Food and Drug Administration. Amplatzer PFO Occluder. Available at: [//www.fda.gov/medicaldevices/productsandmedicalprocedures/deviceapprovalsandclearances/recently-approveddevices/ucm526921.htm](http://www.fda.gov/medicaldevices/productsandmedicalprocedures/deviceapprovalsandclearances/recently-approveddevices/ucm526921.htm). Accessed 01/13/17.
33. Food and Drug Administration. FDA Executive Summary: P120021 AMPLATZER Patent Foramen Ovale (PFO) Occluder. 2016; www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/MedicalDevices/MedicalDevicesAdvisoryCommittee/CirculatorySystemDevicesPanel/UCM502193.pdf. Accessed April, 2017.
34. Food and Drug Administration. PMA P120021: FDA summary of safety and effectiveness data: Patent foramen ovale (PFO) Occluder. 2016; www.accessdata.fda.gov/cdrh_docs/pdf12/P120021B.pdf. Accessed March 19, 2020.
35. Food and Drug Administration. CardioSEAL® Septal Occlusion System with QuikLoad™. Summary of safety and effectiveness data. www.accessdata.fda.gov/cdrh_docs/pdf/P000049b.pdf. Accessed June 15, 2018.
36. Food and Drug Administration. Gore® Cardioform Septal Occluder approval letter. www.accessdata.fda.gov/cdrh_docs/pdf5/P050006s060a.pdf.
37. Food and Drug Administration. Summary of safety and effectiveness data. Gore Cardioform Septal Occluder. March 30, 2018. www.accessdata.fda.gov/cdrh_docs/pdf5/P050006s060b.pdf. Accessed March 30, 2020.
38. Formigari R., DiDonato R.M., et al. Minimally invasive or interventional repair of atrial septal defects in children: experience in 171 cases and comparison with conventional strategies, *Journal of American College of Cardiology*, 2001; 37:6.
39. Furlan AJ, Reisman M, Massaro J, et al. Closure or medical therapy for cryptogenic stroke with patent foramen ovale. *N Engl J Med*. Mar 15 2012; 366(11):991-999.
40. Ghaderian M, Merajie M, Mortezaeian H et al. A mid-term follow-up of the transcatheter closure of perimembranous ventricular septal defects in children using the amplatzer. *J The Univ Heart Ctr* 2015; 10(4): 182-187.
41. Grohmann J, Hohn R, Fleck T, et al. Transcatheter closure of atrial septal defects in children and adolescents: Single-center experience with the GORE septal occluder. *Catheter Cardiovasc Interv*. Mar 24 2014.
42. Gruenstein DH, Ebeid M, Radtke W, et al. Transcatheter closure of patent ductus arteriosus using the Amplatzer™ duct occluder II (ADO II). *Catheter Cardiovasc Interv*. 2017 MAY; 89(6):1118-1128.
43. Hakeem A, Marmagkiolis K, Hacıoglu Y, et al. Safety and efficacy of device closure for patent foramen ovale for secondary prevention of neurological events: Comprehensive systematic review and meta-analysis of randomized controlled trials. *Cardiovasc Revasc Med*. Nov-Dec 2013; 14(6):349-355.
44. Handke M, Harloff A, et al. Patent foramen ovale and cryptogenic stroke in older patients. *NEJM*, November 2007; 357(22): 2262-2268.
45. Harrer JU, Wessels T, et al. Stroke recurrence and its prevention in patients with patent foramen ovale. *Can J Neurol Sci*. 2006 Feb; 33(1):39-47.

46. Homma S., Sacco R.L., et al. Effect of medical treatment in stroke patients with patent foramen ovale: patent foramen ovale in Cryptogenic Stroke Study, *Circulation* 2002; 105(22): 2625-31.
47. Hung J., Ladzbert M.J., Jenkins K.J., et al. Closure of patent foramen ovale or paradoxical emboli: intermediate-term risk of recurrent neurological events following transcatheter device placement, *J Am College Cardio* 2000; 35: 1311-1316.
48. Inglessis I, Elmariah S, Rengifo-Moreno PA, et al. Long-term experience and outcomes with transcatheter closure of patent foramen ovale. *JACC Cardiovasc Interv.* Nov 2013; 6(11):1176-1183.
49. Instruction for Use: Amplatzer® Septal Occluder System. In: AGA Medical Corporation; Golden Valley M, ed.
50. Iverekk H.R., Bone I., Lees K.R. Interatrial septal abnormalities and stroke: a meta-analysis of case-control studies, *Neurology* 2000; 55(8).
51. Jalal ZZ, Hascoët SS, Gronier CC, et al. Long-Term Outcomes After Percutaneous Closure of Ostium Secundum Atrial Septal Defect in the Young: A Nationwide Cohort Study. *JACC Cardiovasc Interv.* 2018 Apr 21;11(8).
52. Javois AJ, Rome JJ, Jones TK, et al. Results of the U.S. Food and Drug Administration continued access clinical trial of the GORE HELEX septal occluder for secundum atrial septal defect. *JACC Cardiovasc Interv.* Aug 2014; 7(8):905-912.
53. Kent DM, Dahabreh IJ, Ruthazer R, et al. Device closure of patent foramen ovale after stroke: pooled analysis of completed randomized trials. *J Am Coll Cardiol.* Mar 01 2016;67(8):907-917.
54. Kernan WN, Ovbiagele B, Black HR, et al. Guidelines for the prevention of stroke in patients with stroke and transient ischemic attack: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke.* Jul 2014; 45(7):2160-2236.
55. Khairy P, O'Donnell CP, Landzberg MJ. Transcatheter closure versus medical therapy of patent foramen ovale and presumed paradoxical thromboemboli: a systematic review. *Ann Intern Med.* 2003 Nov 4; 139(9):753-60.
56. Khan AR, Bin Abdulhak AA, Sheikh MA, et al. Device closure of patent foramen ovale versus medical therapy in cryptogenic stroke: a systematic review and meta-analysis. *JACC Cardiovasc Interv.* Dec 2013; 6(12):1316-1323.
57. Kitsios GD, Dahabreh IJ, Abu Dabrh AM, et al. Patent foramen ovale closure and medical treatments for secondary stroke prevention: a systematic review of observational and randomized evidence. *Stroke.* Feb 2012; 43(2):422-431.
58. Kitsios GD, Thaler DE, Kent DM. Potentially large yet uncertain benefits: a meta-analysis of patent foramen ovale closure trials. *Stroke* 2013; 44(9):2640-3.
59. Kotowycz MA, Therrien J, Ionescu-Ittu R, et al. Long-term outcomes after surgical versus transcatheter closure of atrial septal defects in adults. *JACC Cardiovasc Interv.* May 2013; 6(5):497-503.
60. Kwong JS, Lam YY, Yu CM. Percutaneous closure of patent foramen ovale for cryptogenic stroke: a meta-analysis of randomized controlled trials. *Int J Cardiol.* Oct 9 2013; 168(4):4132-4138.
61. Lansberg MG, O'Donnell MJ, Khatri P, et al. Antithrombotic and thrombolytic therapy for ischemic stroke: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American

- College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest*. Feb 2012;141(2 Suppl):e601S-636S.
62. Late-Breaking Abstracts: 57th Annual Scientific Meeting of the American Headache Society. *Headache: The Journal of Head and Face Pain*. 2015;55:249-265.
 63. Lee PH, Song JK, Kim JS, et al. Cryptogenic Stroke and High-Risk Patent Foramen Ovale: The DEFENSE-PFO Trial. *J Am Coll Cardiol*. May 22 2018; 71(20): 2335-2342.
 64. Li J, Liu J, Liu M, et al. Closure versus medical therapy for preventing recurrent stroke in patients with patent foramen ovale and a history of cryptogenic stroke or transient ischemic attack. *Cochrane Database Syst Rev*. Sep 08 2015(9):CD009938.
 65. Lip PZ, Lip GY. Patent foramen ovale and migraine attacks: a systematic review. *Am J Med*. May 2014; 127(5):411-420.
 66. Mandal KD, Su D, Pang Y. Long-term outcome of transcatheter device closure of perimembranous ventricular septal defects. *Front Pediatr*. 2018 May 3;6:128.
 67. Marshall A.C. and Lock J.E. Valvular and congenital heart disease: structural and compliant anatomy of the patent foramen ovale in patients undergoing transcatheter closure, *American Heart Journal* 2000; 140(2).
 68. Martin F, Sanchez PL, Doherty E, et al. Percutaneous transcatheter closure of patent foramen ovale in patients with paradoxical embolism. *Circulation*. 2002 Aug 27; 106(9):1121-6.
 69. Mas JL, Derumeaux G, Guillon B, et al. Patent foramen ovale closure or anticoagulation vs. antiplatelets after stroke. *N Engl J Med*. Sep 14 2017;377(11):1011-1021.
 70. Mattle HP, Evers S, Hildick-Smith D, et al. Percutaneous closure of patent foramen ovale in migraine with aura, a randomized controlled trial. *Eur Heart J*. Jul 07 2016;37(26):2029-2036.
 71. McDaniel N.L. Ventricular and atrial septal defects, *Pediatrics in Review* 2001; 22(8).
 72. Meier B, Kalesan B, Mattle HP et al. Percutaneous closure of patent foramen ovale in cryptogenic embolism. *N Engl J Med* 2013; 368(12):1083-91.
 73. Messe SR, Gronseth G, Kent DM, et al. Practice advisory: Recurrent stroke with patent foramen ovale (update of practice parameter): Report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology. *Neurology*. Aug 23 2016;87(8):815-821.
 74. Messe SR, Kasner SE. Is closure recommended for patent foramen ovale and cryptogenic stroke? Patent foramen ovale in cryptogenic stroke: not to close. *Circulation*. Nov 4 2008; 118(19):1999-2004.
 75. Messe Sr, Silverman IE, Kizer JR, et al. Practice parameter: recurrent stroke with patent foramen ovale and atrial septal aneurysm: report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology*. 2004 Apr 13; 62(7):1042-50.
 76. Mohr J.P., Thompson J.L.P., et al. A comparison of Warfarin and aspirin for the prevention of recurrent ischemic stroke, *N Engl J Med* 2001; 345(20): 1444-1451.
 77. Mojadidi MK, Gevorgyan R, Nouredin N, et al. The effect of patent foramen ovale closure in patients with platypnea-orthodeoxia syndrome. *Catheter Cardiovasc Interv*. Oct 2015; 86(4):701-707.
 78. Nagaraja V, Raval J, Eslick GD, et al. Is transcatheter closure better than medical therapy for cryptogenic stroke with patent foramen ovale? A meta-analysis of randomized trials. *Heart Lung Circ*. Nov 2013; 22(11):903-909.

79. Nagpal SV, Lerakis S, Flueckiger PB, et al. Long-term outcomes after percutaneous patent foramen ovale closure. *Am J Med Sci*. Sep 2013; 346(3):181-186.
80. National heart, lung, and blood institute. What is patent ductus arteriosus? Available at: www.nhlbi.nih.gov/health/health-topics/topics/pda/. Accessed April 27, 2016.
81. National Institute for Clinical Excellence. Endovascular closure of atrial septal defect. October 2004. Available at: www.nice.org.uk/guidance/ipg96/chapter/1-Guidance.
82. National Institute for Clinical Excellence. Endovascular closure of patent ductus arteriosus. October 2004. Available at: www.nice.org.uk/Guidance/IPG97
83. National Institute for Clinical Excellence. Transcatheter endovascular closure of perimembranous ventricular septal defect. March 2010. Available at: www.nice.org.uk/guidance/ipg336/chapter/1-guidance
84. National Institute for Clinical Excellence. Percutaneous closure of patent foramen ovale to prevent recurrent cerebral embolic events. December 2013. Available at www.nice.org.uk/guidance/ipg472/chapter/1-Recommendations.
85. National Institute for Clinical Excellence. Percutaneous closure of patent foramen ovale for recurrent migraine. December 2010. Available at www.nice.org.uk/guidance/ipg370.
86. Nedeltchev K., Arnold M., et al. Outcome of patient with cryptogenic stroke and patent foramen ovale, *J Neurol Neurosurg Psychiatry* 2002; 27: 347-350.
87. Ntaios G, Papavasileiou V, Makaritsis K, et al. PFO closure vs. medical therapy in cryptogenic stroke or transient ischemic attack: a systematic review and meta-analysis. *Int J Cardiol*. Oct 30 2013; 169(2):101-105.
88. Nyboe C, Hjortdal VE, Nielsen-Kudsk JE. First experiences with the GORE((R)) Septal Occluder in children and adults with atrial septal defects. *Catheter Cardiovasc Interv*. Nov 15 2013; 82(6):929-934.
89. O’Gara PT, Messe SR, et al. Percutaneous device closure of patent foramen ovale for secondary stroke prevention: A call for completion of randomized clinical trials: a science advisory from the American Heart Association/American Stroke Association and the American College of Cardiology Foundation. *Circulation* 2009; 119; 2743-2747; originally published online May 11, 2009; DOI: 10.1161/Circulationaha.109.192272.
90. Oho S, Ishizawa A, et al. Transcatheter closure of atrial septal defects with the Amplatzer septal occluder – a Japanese clinical trial. *Circ J* 2002; 66(9):791-4.
91. Onorato E, Melzi G, et al. Patent foramen ovale with paradoxical embolism: mid-term results of transcatheter closure in 256 patients. *J Interv Cardiol*. 2003 Feb; 16(1):43-50.
92. Paciaroni M, Agnelli G, Bertolini A, et al. Risk of recurrent cerebrovascular events in patients with cryptogenic stroke or transient ischemic attack and patent foramen ovale: the FORI (Foramen Ovale Registro Italiano) study. *Cerebrovasc Dis*. 2011; 31(2):109-116.
93. Pandit A, Aryal MR, Pandit AA, et al. Amplatzer PFO occluder device may prevent recurrent stroke in patients with patent foramen ovale and cryptogenic stroke: a meta-analysis of randomised trials. *Heart Lung Circ*. Apr 2014; 23(4):303-308.
94. Parikh R, Bal N, Pandya D, et al. Comparing outcomes of medical management and percutaneous closure in patients with patent foramen ovale and cryptogenic stroke [abstract]. *Chest*. 2008;134(4):27001.
95. Patti G, Pelliccia F, Gaudio C, et al. Meta-analysis of net long-term benefit of different therapeutic strategies in patients with cryptogenic stroke and patent foramen ovale. *Am J Cardiol*. Mar 15 2015; 115(6):837-843.

96. Pezzini A, Grassi M, Lodigiani C, et al. Propensity Score-Based Analysis of Percutaneous Closure Versus Medical Therapy in Patients With Cryptogenic Stroke and Patent Foramen Ovale: The IPSYS Registry (Italian Project on Stroke in Young Adults). *Circ Cardiovasc Interv.* Sep 2016;9(9).
97. Pickett CA, Villines TC, Ferguson MA, et al. Percutaneous closure versus medical therapy alone for cryptogenic stroke patients with a patent foramen ovale: meta-analysis of randomized controlled trials. *Tex Heart Inst J.* Aug 2014; 41(4):357-367.
98. Pineda AM, Nascimento FO, Yang SC, et al. A meta-analysis of transcatheter closure of patent foramen ovale versus medical therapy for prevention of recurrent thromboembolic events in patients with cryptogenic cerebrovascular events. *Catheter Cardiovasc Interv.* Nov 15 2013; 82(6):968-975.
99. Podnar T., Martanovic P., et al. Morphological variations of secundum-type atrial septal defects: feasibility for percutaneous closure using Amplatzer septal occluders, *Catheterization and Cardiovascular Interventions*, 2001; 53:386-391.
100. Rengifo-Moreno P, Palacios IF, Junpaparp R et al. Patent foramen ovale transcatheter closure vs. medical therapy on recurrent vascular events: a systematic review and meta-analysis of randomized controlled trials. *Eur Heart J* 2013.
101. Rhodes JF, Jr., Goble J. Combined prospective United States clinical study data for the GORE((R)) HELEX((R)) septal occluder device. *Catheter Cardiovasc Interv.* May 1 2014; 83(6):944-952.
102. Rigatelli G, Pedon L, Zecchel R, et al. Long-Term Outcomes and Complications of Intracardiac Echocardiography-Assisted Patent Foramen Ovale Closure in 1,000 Consecutive Patients. *J Interv Cardiol.* Oct 2016;29(5):530-538.
103. Rogers T, Slack M, Waksman R. Overview of the 2016 US Food and Drug Administration Circulatory System Devices Panel Meeting on the Amplatzer Patent Foramen Ovale Occluder. *Am J Cardiol.* Jan 01 2017;119(1):153-155.
104. Sacco RL, Adams R, et al. Guidelines for Prevention of stroke in patients with ischemic stroke or transient ischemic attack: a statement for healthcare professionals from the American Heart Association/American Stroke Association Council on Stroke: co-sponsored by the Council on Cardiovascular Radiology and Intervention. *Stroke* 2006; 37; 577-617.
105. Saver JL, Carroll JD, Thaler DE, et al. Long-term outcomes of patent foramen ovale closure or medical therapy after stroke. *N Engl J Med.* Sep 14 2017;377(11):1022-1032.
106. Shah R, Nayyar M, Jovin IS, et al. Device closure versus medical therapy alone for patent foramen ovale in patients with cryptogenic stroke: a systematic review and meta-analysis. *Ann Intern Med.* Mar 6 2018;168(5):335-342.
107. Slottow TL, Steinberg DH, Waksman R. Overview of the 2007 Food and Drug Administration Circulatory System Devices Panel meeting on patent foramen ovale closure devices. *Circulation.* Aug 7 2007; 116(6):677-682.
108. Sndergaard L, Kasner SE, Rhodes JF, et al. Patent foramen ovale closure or antiplatelet therapy for cryptogenic stroke. *N Engl J Med.* Sep 14 2017;377(11):1033-1042.
109. Snijder RJ, Luermans JG, de Heij AH, et al. Patent Foramen Ovale With Atrial Septal Aneurysm Is Strongly Associated With Migraine With Aura: A Large Observational Study. *J Am Heart Assoc.* Dec 01 2016;5(12).
110. Søndergaard L, Kasner SE, Rhodes JF, et al. Patent foramen ovale closure or antiplatelet therapy for cryptogenic stroke. *N Engl J Med.* Sep 14 2017;377(11):1033-1042.

111. Stortecky S, da Costa BR, Mattle HP, et al. Percutaneous closure of patent foramen ovale in patients with cryptogenic embolism: a network meta-analysis. *Eur Heart J*. Jan 7 2015; 36(2):120-128.
112. Stout KK, Daniels CJ, Abouhosn JA, et al. 2018 AHA/ACC guideline for the management of adults with congenital heart disease. *Circulation*. Apr 2 2019; 139: e698 –e800.
113. Suchon E, Pieculewicz M, et al. Transcatheter closure as an alternative and equivalent method to the surgical treatment of atrial septal defect in adults: comparison of early and late results. *Med Sci Monit* 2009; 15(12):CR612-7.
114. Taeed R., Shim D., et al. One-year follow-up of the Amplatzer device to close atrial septal defects, *American Journal of Cardiology*, 2001; 84:1.
115. The AMPLATZER® septal occluder and delivery system, AGA Medical Corporation, Golden Valley, Minnesota.
116. Thomson JD, Hildick-Smith D, Clift P, et al. Patent foramen ovale closure with the gore septal occluder: initial UK experience. *Catheter Cardiovasc Interv*. Feb 15 2014; 83(3):467-473.
117. Tobis JM, Charles A, Silberstein SD, et al. Percutaneous closure of patent foramen ovale in patients with migraine: the PREMIUM Trial. *J Am Coll Cardiol*. Dec 5 2017;70(22):2766-2774.
118. Tobis J, Shenoda M. Percutaneous Treatment of Patent Foramen Ovale and Atrial Septal Defects. *J Am Coll Cardiol*. 2012; 60(18):1722-1732.
119. Udell JA, Opatowsky AR, Khairy P, et al. Patent foramen ovale closure vs medical therapy for stroke prevention: meta-analysis of randomized trials and review of heterogeneity in meta-analyses. *Can J Cardiol*. Oct 2014; 30(10):1216-1224.
120. Wahl A, Juni P, Mono ML, et al. Long-term propensity score-matched comparison of percutaneous closure of patent foramen ovale with medical treatment after paradoxical embolism. *Circulation*. Feb 14 2012; 125(6):803-812.
121. Wallenborn J, Bertog SC, Franke J, et al. Recurrent events after percutaneous closure of patent foramen ovale. *Catheter Cardiovasc Interv*. Oct 1 2013; 82(4):541-546.
122. Warnes CA, Williams RG, et al. ACC/AHA 2008 guidelines for the management of adults with congenital heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Develop Guidelines on the Management of Adults with Congenital Heart Disease). *Circulation* 2008; 118(23):e714-833.
123. Wein T.H., Bornstein N.M. Stroke prevention: cardiac and carotid-related stroke, *Neurologic Clinics* 2000; 18(2).
124. Whitlock RP, Sun JC, Fries SE, et al. Antithrombotic and thrombolytic therapy for valvular disease: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest*. Feb 2012; 141(2 Suppl):e576S-600S.
125. Windecker S, Wahl A, et al. Comparison of medical treatment with percutaneous closure of patent foramen ovale in patients with cryptogenic stroke. *J Am Coll Cardiol*. 2004 Aug 18; 44(4):750-8.
126. Windecker S., Wahl A., et al. Percutaneous closure of patent foramen ovale in patients with paradoxical embolism, *Circulation* 2000; 101:893-898.

127. Wintzer-Wehekind J, Alperi A, Houde C, et al. Long-Term Follow-Up After Closure of Patent Foramen Ovale in Patients With Cryptogenic Embolism. *J Am Coll Cardiol*. Jan 29 2019; 73(3): 278-287.
128. Wohrle J. Closure of patent foramen ovale after cryptogenic stroke. *Lancet*. 2006 Jul 29; 368(9533):350-2.
129. Wolfrum M, Froehlich GM, Knapp G, et al. Stroke prevention by percutaneous closure of patent foramen ovale: a systematic review and meta-analysis. *Heart*. Mar 2014; 100(5):389-395.
130. Xu XD, Liu SX, Zhao XX, et al. Comparison of Medium-term Results of Transcatheter Correction Versus Surgical Treatment for Secundum Type Atrial Septal Defect Combined With Pulmonary Valve Stenosis. *Int Heart J*. Jul 10 2014; 55(4):326-330.
131. Yang J, Lifang Y, Shiqiang Y et al. Transcatheter versus surgical closure of perimembranous ventricular septal defects in children. *Journal of the American College of Cardiology*. 2014 Nov; 63(12):1159-1168.
132. Yang J, Yang L, Wan Y et al. Transcatheter device closure of perimembranous ventricular septal defects: mid-term outcomes. *Eur Heart J*. 2010 Sep; 31(18): 2238-45.
133. Yilmazer MM, Guven B, Vupa-Cilengiroglu O, et al. Improvement in cardiac structure and functions early after transcatheter closure of secundum atrial septal defect in children and adolescents. *Turk J Pediatr*. Jul-Aug 2013; 55(4):401-410.

POLICY HISTORY:

Adopted for Blue Advantage, September 2011

Available for comment September 22 through November 7, 2011

Medical Policy Group, December 2011

Medical Policy Group, October 2013

Medical Policy Group, December 2013

Medical Policy Group, October 2014

Medical Policy Group, July 2015

Medical Policy Group, December 2015

Medical Policy Group, June 2016

Available for comment June 29 through August 13, 2016

Medical Policy Group, January 2017

Medical Policy Group, April 2017

Medical Policy Group, May 2017

Medical Policy Group, June 2018

Medical Policy Group, June 2019

Medical Policy Group, June 2020: Deleted policy section for dates of service on or after April 12, 2011, and prior to August 14, 2016.

This medical policy is not an authorization, certification, explanation of benefits, or a contract. Eligibility and benefits are determined on a case-by-case basis according to the terms of the member's plan in effect as of the date services are rendered. All medical policies are based on (i) research of current medical literature and (ii) review of common medical practices in the treatment and diagnosis of disease as of the date hereof. Physicians and other providers are solely responsible for all aspects of medical care and treatment, including the type, quality, and levels of care and treatment.

This policy is intended to be used for adjudication of claims (including pre-admission certification, pre-determinations, and pre-procedure review) in Blue Cross and Blue Shield's administration of plan contracts.